

Effect of Pre-Harvest Treatments and Bagging for Quality Improvement of Guava (*Psidium guajava* L.) cv L- 49 Fruits

Saransh Saxena, Priyamvada Sonkar, Nidhi Patel,
Jyoti Singh, Jitendra Bhandari

Received 2 April 2021, Accepted 19 June 2021, Published on 8 June 2022

ABSTRACT

The investigation was carried out at Instructional cum Research Fruit Orchard, Department of Fruit Science, KNK. College of Horticulture, Mandsaur (MP), during November 2019 to January 2020. The treatments comprised of pre harvest spraying of CaCl₂, Ascorbic acid and bagging with Brown paper and yellow color polythene to study with quality improvement of guava fruit. The experiment was laid out in Randomized Complete Block Design with three replications. Treatment CaCl₂ @ 2% with Brown paper (T₁₁) fruit bagging showed significant effects on different parameters studied. It was observed that the maximum acidity (0.57%) was recorded in treatment T₀ (control), while maximum total soluble solids (°B) (11.50), TSS: Acidity ratio (26.63), Ascorbic acid content (192.39 mg/100g pulp), pectin content (1.05

%), pH (4.73), Chlorophyll content in leaves (46.10 SPAD value) were recorded with the treatment CaCl₂ @ 2% with Brown paper (T₁₁).

Keywords Calcium chloride, Ascorbic acid, Brown paper, Yellow polythene bag, Quality.

INTRODUCTION

Guava (*Psidium guajava* L.) is one of the important fruit cultivated in several tropical and subtropical countries of the world (Pathak *et al.* 2007), it belongs to the family Myrtaceae, originated in Tropical America. It is also known as 'Apple of the tropics' or 'Poor man's apple'. Due to hardy nature of the guava fruit plant, it can withstand upto adverse climatic conditions and grows under a wide range of soil types from sandy loam to clay loam (Dhaliwal and Singla 2002). In India, Guava is the fourth most important fruit crop in area and production after mango, banana and citrus with production of 42.36 lakh tonnes from an area of 2.76 lakh ha (NHB Database 2018). It is an important fruit crop cultivated all over India, especially in the regions of Uttar Pradesh, Madhya Pradesh, Bihar, Maharashtra, West Bengal, Punjab, Chhattisgarh, Haryana, Karnataka and Gujarat.

Saransh Saxena*, Nidhi Patel, Jyoti Singh
Department of Fruit Science, RVSKVV, College of Horticulture,
Mandsaur (MP) 458001, India

Priyamvada Sonkar
Assistant Professor, Department of Fruit Science, RVSKVV,
College of Horticulture, Mandsaur (MP), India

Jitendra Bhandari
Assistant Professor, Department of Horticulture, Mandsaur Uni-
versity, Mandsaur (MP), India
Email: saranshsaxena220@gmail.com

*Corresponding author

Guava fruit of variety Lucknow-49 is a selection from open pollinated population of Allahabad Safeda cultivar from Ganeshkhind, Pune (MH) and is also called as Sardar Guava. Trees are vigorous. The leaves are elliptical-ovate to oblong in shape. Fruits are roundish ovate in shape. Skin color is primrose-yellow. The pulp is whitish, very sweet and tasty with a TSS of 9.5 °B and vitamin C content 130 mg/100 g. It has been observed that pectin content is higher in winter guava than the monsoon fruits (Salunke and Kadam 1995).

Pre-harvest calcium spray is one of the most important practices of new strategies applied in the integrated fruit production systems, improving fruit characteristics to minimize fungicides sprays towards the end of the harvest period, which in turn improves fruit resistance to brown rot (Conway *et al.* 1994). Calcium spray during fruit development provides a safe mode of supplementing endogenous calcium to fresh fruits (Raese and Drake 2000). Application of Ascorbic acid had many stimulating effects on growth and physiological activities of various plants (Abdou *et al.* 2015). Ascorbic acid is a good antioxidant that keeps fruit from darkening and improves destruction of bacteria. Antioxidants are used instead of auxins for improving fruit growth, development and fruiting of trees (Maksoud *et al.* 2009).

Fruit bagging decreases the defects generated due to diseases and insects, and increased flesh firmness and flavor. The most important role of fruit bagging was to effectively protect fruits from physiological factors which led to the significant decrease of the total damaged, degenerative and defective fruits (13.7-33.3%), as compared with non-bagging fruits. Bagging is a physical protection technique used extensively in several fruit crops to improve skin color which also reduce the incidence of disease, insect-pests, mechanical damage, agrochemical residues on the fruit and bird damage (Xu *et al.* 2010).

MATERIALS AND METHODS

The present investigation was conducted during the year 2019-2020 at the Instructional Cum Fruit Research Orchard, Department of Fruit Science, KNK College of Horticulture Mandsaur (MP). Sin-

gle spraying of CaCl₂ and Ascorbic acid were done at 30 days before harvesting of the fruits. Bagging of fruits with Yellow polyethylene bags and Brown paper bags were done one month before harvesting of the fruits of guava fruits comprised of 21 treatments included control. Without spray and no fruit bagging (open fruit) was treated as control. The experiment was conducted in Randomized Complete Block Design (RCBD) with three replications. The fruits were wrapped with respective bagging materials as per the treatments. Single tree was considered as an experimental unit. Hand refractometer was used for determination of TSS in °Brix. Acidity was estimated by simple acid-alkali titration method as described in AOAC in (1970). Value of Total Soluble Solids was divided by acidity percent to obtain TSS: Acidity ratio. Ascorbic acid was estimated by Assay method given by Ranganna (2013). Estimation of pectin according to the methods of Kertesz (1951). The pH of fruit was determined by method suggested by Piper (1966). Chlorophyll content in leaves was estimated by using instrument SPAD-502 chlorophyll meter.

RESULTS AND DISCUSSION

Present study of pre-harvest treatments and bagging have considerable enhancement on the biochemical parameters of guava and the data regarding this is given in Table 1. Application of CaCl₂ and Ascorbic acid along with bagging improves the fruit quality. Their application alone or in combination influenced significantly the chemical constituents of the fruit viz. TSS, ascorbic acid, TSS: acidity ratio, pectin content, pH and chlorophyll content in leaves over the control. The maximum acidity (0.57%) was in treatment T₀ (control), while maximum Total Soluble Solids (11.50 °B), maximum ascorbic acid content (192.39 mg), maximum TSS: Acidity ratio (26.63), maximum pectin content (1.05%), maximum pH (4.73) and maximum chlorophyll content in leaves (46.10) with the application of T₁₁ (CaCl₂ @ 2% with Brown paper) at harvest were significantly superior to (control) T₀.

The improvement in various chemical characteristics is due to application of chemical combination treatment beneficially affected in increasing the various constituents of fruits hence quality improvement

Table 1. Effect of pre-harvest treatments and bagging on quality parameters of guava fruit.

Treatments	TSS (^o B)	Acidity	TSS : Acidity ratio	Ascorbic acid (mg/ 100g pulp)	Pectin (%)	pH	Chloro- phyll (SPAD)
T ₀ Control	8.61	0.57	15.22	149.33	0.76	3.55	30.20
T ₁ CaCl ₂ @ 1%	9.53	0.40	17.67	157.22	0.82	3.92	39.20
T ₂ CaCl ₂ @ 1.5%	9.53	0.43	21.55	164.04	0.91	4.01	39.57
T ₃ CaCl ₂ @ 2%	10.62	0.47	24.35	168.30	0.93	4.04	39.77
T ₄ Ascorbic acid @ 200ppm	9.78	0.45	19.20	148.57	0.78	3.79	37.90
T ₅ Ascorbic acid @ 300 ppm	8.69	0.43	19.74	146.53	0.85	4.01	36.57
T ₆ Ascorbic acid @ 400 ppm	9.61	0.44	21.07	154.05	0.85	4.05	39.03
T ₇ Bagging with Brown paper	10.33	0.46	21.43	191.45	1.02	4.07	41.33
T ₈ Bagging with Yellow polyethylene	10.00	0.49	20.49	183.84	0.97	3.88	39.13
T ₉ CaCl ₂ @ 1% with Brown paper	10.39	0.33	22.47	166.85	0.99	4.10	44.47
T ₁₀ CaCl ₂ @ 1.5% with Brown paper	10.29	0.43	25.84	187.74	0.99	4.15	41.27
T ₁₁ CaCl ₂ @ 2% with Brown paper	11.50	0.47	26.63	192.39	1.05	4.73	46.10
T ₁₂ Ascorbic acid @ 200 ppm with Brown paper	10.28	0.47	20.77	182.76	0.99	4.38	35.50
T ₁₃ Ascorbic acid @ 300 ppm with Brown paper	10.22	0.46	26.33	188.28	0.92	3.88	35.17
T ₁₄ Ascorbic acid @ 400 ppm with Brown paper	9.28	0.43	24.78	159.00	0.97	4.05	37.03
T ₁₅ CaCl ₂ @ 1% with Yellow polyethylene	9.66	0.48	23.79	154.75	1.03	4.05	33.50
T ₁₆ CaCl ₂ @ 1.5% with Yellow polyethylene	10.50	0.41	26.23	180.13	0.91	4.07	36.70
T ₁₇ CaCl ₂ @ 2% with Yellow polyethylene	10.22	0.37	26.47	188.16	0.97	4.11	36.13
T ₁₈ Ascorbic acid @ 200 ppm with Yellow polyethylene	9.62	0.41	23.72	172.59	0.95	4.00	38.27
T ₁₉ Ascorbic acid @ 300 ppm with Yellow polyethylene	9.61	0.41	23.53	163.63	0.93	4.12	36.17
T ₂₀ Ascorbic acid @ 400 ppm with Yellow polyethylene	9.70	0.41	24.55	162.79	0.92	4.07	35.97
SEm ±	0.24	0.02	0.89	3.31	0.03	0.08	0.67
CD at 5%	0.68	0.06	2.54	9.47	0.08	0.23	1.91

reflected in chemical characters of fruit. Guava responds well to the application of calcium, hence CaCl₂ application improves, fruit character and chemical composition through rapid transformation of plant nutrients along with Ascorbic acid application of different concentrations which has stimulating effects over quality parameters of the fruit. Similar findings were also reported by many scientists. Same time in bagging (Brown paper and Yellow polyethylene) enhanced much better chemical composition with the induction of growth hormones, due to which sufficient movement of water and nutrients which finally responsible for growth and development in qualitative way of fruits and other bio-chemical characters. The present findings are in accordance with the results reported by Rahman *et al.* (2018) and Islam *et al.* (2019).

CONCLUSION

All the pre harvest treatments were found better than control in improving fruit quality while the pre

harvest treatment of CaCl₂ @ 2% with Brown paper was found superior to increase the fruit quality than all other treatments. This treatment was found to have very low spots and no infestation. Hence it should be practiced in guava crop to produce fruits with good quality.

REFERENCES

- Abdou MAH, Ahmed EET, Ibrahim TIE (2015) Effect of nitrogen sources and Ascorbic acid on growth and essential oil production of geranium (*Pelargonium graveolens* L.) plants. *Sci J Flowers Ornamental Pl* 2(2): 157-165.
- AOAC (2000) Official Methods of Analysis. Association of the Official Analytical Chemists, Washington, DC 13th edn.
- Conway WS, Sams CE, Kelman A (1994) Enhancing the natural resistance of plant tissue to post-harvest diseases through calcium applications. *Hort Sci* 29: 751-761.
- Dhaliwal GS, Singla R (2002) Studies on the time of anthesis and dehiscence in different genotypes of guava in winter and rainy season crops. *Ind J Hort* 59: 157-161.
- Islam MT, Shamsuzzoha M, Rahman MS, Bari MA, Akter MM, Khatun A, Huque R, Uddin MS (2019) Effect of bagging time on fruit quality and shelf life of Mango (*Mangifera indica* L.) cv. Langra in Bangladesh. *Int J Agric Environ Biores* 4: 279-289.

- Kerteszi ZI (1951) *The Pectic Substances*. Interscience Publishers, New York.
- Maksoud MA, Salah MS, El-Shamma, Faud AA (2009) The effect of biofertilizers and antioxidants on olive trees under calcareous soil conditions. *World J Agric Sci* 5: 350-352.
- NHB (2018). National Horticulture Database. National Horticulture Board. Govt India, Gurgaon, India. www.nhb.gov.in.
- Pathak RK, Singh G, Kishun R, Chandra R (2007) Improvement of Guava (*Psidium guajava* L.). *Through Breed* 85(1): 0567-7572 .
- Piper CS (1966) *Soil and Plant analysis*. Hans Publisher, Bombay.
- Raese JT, Drake SR (2000) Effect of calcium sprays, time of harvest, cold storage, and ripeness on fruit quality of 'Anjou' Pears. *J Pl Nutr* 23: 843–853.
- Rahman MM, Hossain MM, Rahim MA, Rubelmd HK, Islam MZ (2018) Effect of pre-harvest fruit bagging on post-harvest quality of Guava cv Swarupkathi. *Fundam Appl Agric* 3 (1): 363–371.
- Rangana S (2013) *Manual for analysis of fruit and vegetable products*. Tata Mcgraw Co Pvt Ltd New Delhi.
- Salunkhe DK, Kadam SS (1995) *Handbook of fruit science and technology production, composition, storage and processing*. Marcel Dekker, Inc., pp 419-433.
- Xu HX, Chen JW, Xie M (2010) Effect of different light transmittance paper bags on fruit quality and anti-oxidant capacity in Loquat. *J Sci Food Agric* 90: 1783–1788.