

Effect of Sewage Water Irrigation on Micronutrient and Toxic Metal Contents in Plants

Deepak Kumar, S. K. Singh* , Sardul Mann, Sandeep Bhakar

Received 14 June 2020, Accepted 4 September 2020, Published on 5 October 2020

ABSTRACT

A survey was conducted in Kaithal, Kurukshetra, Karnal, Panipat and Faridabad cities of South-Eastern Haryana State, from where plant samples of different vegetables, forage and field crops collected from sewer water irrigated field and analyzed for micronutrient and toxic metal contents in plants at Department of Soil Science, CCS Haryana Agricultural University, Hisar. It was observed from the data that the copper concentration in different plant samples varied from 7.0-70.3 mg kg⁻¹ and 7.5-63.5 mg kg⁻¹ whereas iron content ranged from 90.0-1049.0 mg kg⁻¹ and 464.0-1287.0 mg kg⁻¹ in shoots and roots, respectively. Zinc concentration ranged from 6.0-140.3 mg kg⁻¹ and 32.5-199.9 mg kg⁻¹ whereas manganese content ranged from 18.7-106.9 mg kg⁻¹ and 31.6-288.7 mg kg⁻¹ in shoots and roots, respectively. The cadmium content in shoots and roots of sewage water irrigated plants ranged from 2.5-18.0 mg kg⁻¹ and 4.5-9.0 mg

kg⁻¹ respectively whereas lead content in different plant samples was found to vary from 4.9-68.9 mg kg⁻¹ in shoots and 12.0-60.9 mg kg⁻¹ in roots. The nickel content ranged from 1.8-9.6 and 3.4-22.4 mg kg⁻¹ whereas chromium content ranged from 0.4-3.9 mg kg⁻¹ and 0.5-10.4 mg kg⁻¹ in shoots and roots, respectively. It was also observed that the concentration of heavy metals except Cd and Pb in sewage water irrigated plant samples were lower than maximum phytotoxicity limit in plants (20-100, 150-200, >500, >500, >0.8, >20, >50 and >10 mg kg⁻¹ for Cu, Zn, Fe, Mn, Cd, Pb, Ni and Cr, respectively). The concentration of the micronutrients except Cu and all the toxic metals were found to be more in roots than in shoots.

Keywords: Vegetables, Forage , Field crops, Micronutrient, Toxic metal.

INTRODUCTION

Agricultural lands particularly near urban areas are irrigated with industrial, city or municipal waste water due to low availability of fresh water, undemanding accessibility and discarding problems of waste water. In general, this resource contains substantial amounts of beneficial nutrients and toxic pollutants, which are creating opportunities and problems for agricultural production, respectively (Alghobar and Suresha 2017). Heavy metals are injurious to humans and animals due to their solubility in water, non-biodegradable nature and their potential to build up in var-

Deepak Kumar, S. K. Singh*, Sardul Mann, Sandeep Bhakar
Chaudhary Charan Singh Haryana Agricultural University,
Krishi Vigyan Kendra, Fatehabad, Haryana, India
Email :sksinghphd@gmail.com

*Corresponding author

ious body parts. These are even toxic and detrimental to humans at very low concentrations as there is no proper route for their removal from the body. Waste water contains large quantity of metals such as lead (Pb), nickel (Ni), iron (Fe), copper (Cu), cadmium (Cd), chromium (Cr), zinc (Zn) (Nazir et al. 2015). On the other hand, soils irrigated with waste water are beneficial for farmers as these have enhanced organic matter content and essential plant nutrients as compared to fresh water irrigated soils and impart to lessen the expenditures of fertilizers (Rai et al. 2011) mainly in soils situated near industries and peri-urban areas. This issue needs special attention, when agricultural lands are continuously irrigated with untreated sewage water for longer times to cultivate crops/vegetables. Heavy metal bioaccumulation in the food chain can be highly dangerous to human health (Dadar et al. 2016). Under wheat crop, sewage irrigation for four decades resulted into a significant buildup of zinc (141 %), copper (219 %), iron (514 %), nickel (75.0 %), and lead (28.1 %) in sewage-irrigated soils over adjacent tube well water-irrigated ones (Meena et al. 2016). Concentration of heavy metals in sugar beet plant grown on industrial effluent was higher as compared to their tolerance level indicating their accumulation in plants (Rajput et al. 2017). Thus, the survey was conducted to study the effect of sewage water irrigation on micronutrient and toxic metal contents in plants primarily to quantify the concentration of metals in irrigation waste water crops/vegetables cultivated in Kaithal, Kurukshetra, Karnal, Panipat and Faridabad cities of South-Eastern Haryana State.

MATERIALS AND METHODS

Plant samples of different vegetables, forage and field crops were collected from sewer water irrigated field of Kaithal, Kurukshetra, Karnal, Panipat and Faridabad cities of Haryana State. The plant samples collected were first washed with tap water to remove the adhering soil particles, followed by acidified de-ionized water and then with double distilled water. The excess water was removed by gentle shaking and pressing against the filter paper. The washed plant samples were first air dried and then put in the paper bags and placed it in an air draft oven at $65\pm 2^{\circ}\text{C}$ to a constant weight. Thereafter, the shoots and roots were separated and ground in a stainless steel grinder and

stored in small paper bags for laboratory analysis. In order to determine the micronutrients and heavy metals viz. Cu, Zn, Fe, Mn, Cd, Pb, Ni and Cr in plant parts (shoot and root), 0.5 g of ground and well mixed material was digested in a di-acid mixture of nitric acid and perchloric acid (4:1). After digestion the volume was made to 50 ml with distilled water, filtered and stored in well washed plastic bottles. The Cu, Zn, Fe, Mn, Cd, Pb, Ni and Cr were estimated from the aliquot by using the Atomic Absorption Spectrophotometer.

RESULTS AND DISCUSSION

Micronutrient concentration

The data presented in Table 1 showed that the copper concentration in different plant samples collected from Kaithal, Kurukshetra, Karnal, Panipat and Faridabad cities receiving sewage water irrigation varied from 7.0-70.3 mg kg⁻¹ and 7.5-63.5 mg kg⁻¹ in shoots and roots respectively. The concentration of Cu in shoot samples was maximum (70.3 mg kg⁻¹) in tomato (*Lycopersicon esculantum*) and minimum (7.0 mg kg⁻¹) was found in cauliflower (*Brassica oleracea* var. botrytis). While, in case of root samples the Cu concentration was maximum (63.5 mg kg⁻¹) in brinjal (*Solanum melongena*) and minimum (7.5 mg kg⁻¹) in onion (*Allium cepa*). As per the maximum permissible limit of Cu 4.0 to 15.0 mg kg⁻¹ for most plant species (Alloway, 1968), all of the plant samples in the present study were found below the maximum phytotoxicity level of 20-100 mg kg⁻¹ as mentioned by Mortvedt et al. (1991), Tandon (1995). Anwar et al. (2016) reported that the maximum Cu concentrations were accumulated in mint roots as compared to coriander and fenugreek. The results are in confirmation with the findings of (Alawsy et al. 2018, Chaoua et al. 2019). Zinc concentration in plants ranged from 6.0-140.3 mg kg⁻¹ and 32.5-199.9 mg kg⁻¹ in shoot and root respectively (Table 1). In case of shoot samples the Zn concentration was maximum (140.3 mg kg⁻¹) in Jowar (*Sorghum bicolor*) and minimum (6.0 mg kg⁻¹) in Okra (*Ablemoschus esculantus*). Whereas, in roots samples it was noticed maximum (199.9 mg kg⁻¹) in tomato (*Lycopersicon esculantum*) and minimum (32.5 mg kg⁻¹) in onion (*Allium cepa*). The content of Zn in all the plant samples was below the maximum phytotoxicity levels of 150-200

mg kg⁻¹. The biomass of all the vegetables were negatively affected when irrigated with sewage water. Zinc was accumulated in the sequence of leaves > roots > shoots under polluted water irrigation (Anwar et al. 2016). Results are consistent with the findings of (Alawsy et al. 2018, Chaoua et al. 2019). The iron content in shoots and roots of plant (Table 1) ranged from 90.0-1049.0 mg kg⁻¹ and 464.0-1287.0 mg kg⁻¹ respectively. In shoots, it was maximum (1049.0 mg kg⁻¹) in brinjal (*Solanum melongena*) and minimum (90.0 mg kg⁻¹) in okra (*Ablemoschus esculantus*). Likewise in roots, the concentration was maximum (1287.0 mg kg⁻¹) in ridge gourd (*Lijaria sinaria*) and minimum (464.0 mg kg⁻¹) in okra (*Ablemoschus esculantus*) plants. As the critical limit of phytotoxicity reported by Mortvedt et al. (1991), Tandon (1995) for Fe is > 500 mg kg⁻¹, hence the level of Fe content in different plant parts of the study area was much above the critical levels. The results are in confirmation with the results of (Naz et al. 2018, Alawsy et al. 2018). The shoots and roots manganese content of different plants collected from the study area ranged from 18.7-106.9 mg kg⁻¹ and 31.6-288.7 mg kg⁻¹ respectively (Table 1). The concentration of Mn was maximum (106.9 mg kg⁻¹) in spinach (*Beta vulgaris*) leaf and minimum (18.7 mg kg⁻¹) in coriander (*Coriandrum sativum* L.) leaves. While in roots the concentration of Mn was highest (288.7 mg kg⁻¹) in barseem (*Alexandrum trifolium*) and lowest (31.6 mg kg⁻¹) in kidney bean (*Phaseolus vulgaris* L.). The Mn content in different plant samples were much below the phytotoxicity level of > 500 mg kg⁻¹ as mentioned by Mortvedt et al. (1991), Tandon (1995). The low content of Mn in plants can be attributed to the antagonistic effect of toxic metal present in the sewage (Knezek and Greinert 1971). Alawsy et al. (2018) found that the absorbed Mn by the maize and barley crops is increased with increasing concentration of Mn in sewage water.

Toxic metal concentration

The cadmium content in shoots and roots of sewage water irrigated plants ranged from 2.5-18.0 mg kg⁻¹ and 4.5-9.0 mg kg⁻¹ respectively (Table 1). The content in shoot samples was maximum (18.0 mg kg⁻¹) in bitter melon (*Momordica charantia*) and minimum

(2.5 mg kg⁻¹) in water melon (*Citrullus vulgaris*). In root samples the Cd content was maximum (9.0 mg kg⁻¹) in tomato (*Lycopersicon esculantum*) and minimum (4.5 mg kg⁻¹) in chilli (*Capsium annum*). The Cd content in all of the plant samples was above the toxicity thresholds limit of 0.8 mg kg⁻¹ as mentioned by Alloway (1968). Reported that mean concentrations of Cadmium (Cd) in plant samples of Palak, Amaranthus, Spinach, Coriander, Green chillies and Faragrass under sewage irrigated condition were higher as compared to ground water irrigation and the corresponding values were relatively lower under no irrigation (control). Naz et al. (2018) also found that the vegetables irrigated with sewage water had greater concentrations of Cd which was higher than the safe limits recommended by WHO (1996). The results are in confirmation with the results of (Khaled et al. 2016, Chaoua et al. 2019). The lead content in different plant samples was found to vary from 4.9-68.9 mg kg⁻¹ in shoots and 12.0-60.9 mg kg⁻¹ in roots (Table 1). It was maximum in shoot (68.9 mg kg⁻¹) in brinjal (*Solanum melongena*) and roots (60.9 mg kg⁻¹) in onion (*Allium cepa*). Whereas, it's minimum concentration was found in shoots (4.9 mg kg⁻¹) of okra (*Ablemoschus esculantus*) and roots (12.0 mg kg⁻¹) of kidney bean (*Phaseolus vulgaris* L.). The Pb content in most of the plants under study was above the critical limit of phytotoxicity of > 20 mg kg⁻¹ as mentioned by Mortvedt et al. (1991), Tandon (1995) and hence proper monitoring of sewage water is essential before its use for irrigation in the present study area. Anwar et al. (2016) reported that the maximum Pb concentrations were accumulated in mint roots as compared to coriander and fenugreek. The vegetables irrigated with sewage water had greater concentrations of Pb which was higher than the safe limit (Naz et al. 2018). The results are in consistent with the findings of (Alawsy et al. 2018, Chaoua et al. 2019). The nickel content in various sewage water irrigated plants ranged from 1.8-9.6 and 3.4-22.4 mg kg⁻¹ in shoots and roots respectively (Table 1). The concentration was maximum (9.6 mg kg⁻¹) in shoots of spinach (*Beta vulgaris*) and minimum (1.8 mg kg⁻¹) in okra (*Ablemoschus esculantus*). In roots samples, maximum Ni concentration (22.0 mg kg⁻¹) was noticed in ridge gourd (*Lajria sinaria*) and minimum (3.4 mg kg⁻¹) in kidney bean (*Phaseolus vulgaris* L.).

As the critical limit of Ni phytotoxicity for plant tissue is $> 50 \text{ mg kg}^{-1}$ as mentioned by Mortvedt et al. (1991), Tandon (1995), so all the plants are safe for consumption in the study area as far as Ni content is concerned. Wheat roots accumulated significantly higher amount of Ni in sewage-irrigated land compared to bore well-irrigated land. The same trend was noticed in stem. In general, concentration of Ni was higher in root followed by stem and lower in grain (Salakinkop and Hunshal 2014). The results are in confirmation with the results of (Salakinkop and Hunshal 2014, Naz et al. 2018). The chromium content in shoots and roots of different plants (Table 1) was found to range from $0.4\text{-}3.9 \text{ mg kg}^{-1}$ and $0.5\text{-}10.4 \text{ mg kg}^{-1}$ respectively. Maximum Cr content (3.9 mg kg^{-1}) was found in brinjal shoot (*Solanum melongena*) and minimum (0.4 mg kg^{-1}) in coriander (*Coriandrum sativum* L.) and round melon (*Citrullus vulgaris* var. *Fistulosus*). Similarly in roots also the concentration was more (10.4 mg kg^{-1}) in barseem (*Alexandrum trifolium*) and less (0.5 mg kg^{-1}) in okra (*Ablemoschus esculantus*). The Cr content in all the above ground plant parts in the study area was below permissible level of 10 mg kg^{-1} (Alloway 1968) in all the crop except barseem (Alexandrum

trifolium) and hence at present there is no toxicity of Cr in crop grown of sewage water irrigated area in different districts of Haryana under study. The results of the present investigation are in line with the finding of Salakinkop and Hunshal (2014), Naz et al. (2018).

The chromium content in shoots and roots of different plants (Table 1) was found to range from $0.4\text{-}3.9 \text{ mg kg}^{-1}$ and $0.5\text{-}10.4 \text{ mg kg}^{-1}$ respectively. Maximum Cr content (3.9 mg kg^{-1}) was found in brinjal shoot (*Solanum melongena*) and minimum (0.4 mg kg^{-1}) in coriander (*Coriandrum sativum* L.) and round melon (*Citrullus vulgaris* var. *Fistulosus*). Similarly in roots also the concentration was more (10.4 mg kg^{-1}) in barseem (*Alexandrum trifolium*) and less (0.5 mg kg^{-1}) in okra (*Ablemoschus esculantus*). The Cr content in all the above ground plant parts in the study area was below permissible level of 10 mg kg^{-1} (Alloway 1968) in all the crop except barseem (*Alexandrum trifolium*) and hence at present there is no toxicity of Cr in crop grown of sewage water irrigated area in different districts of Haryana under study. The results of the present investigation are in line with the finding of Salakinkop and Hunshal (2014), Naz et al. (2018).

Table 1. Effect of sewage water irrigation on micronutrients and toxic metals contents (mg kg^{-1}) in different field crops.

Crops	Parts	Kaithal							
		Cu	Zn	Fe	Mn	Cd	Pb	Ni	Cr
Okra (<i>Ablemoschus esculantus</i>)	Shoot	46.3	106.5	830	44.6	7.0	19.8	3.2	1.9
	Root	24.8	106.8	565	92.7	5.5	26.0	3.6	3.8
Bitter gourd (<i>Momordica charantia</i>)	Shoot	34.8	45.5	571	19.9	4.0	17.1	3.6	ND
	Root	10.3	129.9	542	73.1	8.5	22.1	6.4	1.6
Mint (<i>Monarda arvensis</i>)	Shoot	69.3	53.5	850	70.4	4.0	14.5	5.5	3.7
	Root	39.8	150.0	1191	110.8	5.0	20.3	83	6.6
Chilli (<i>Capsicum annum</i>)	Shoot	17.3	116.5	745	53.3	5.0	12.6	3.7	0.5
	Root	54.3	121.9	1189	100.6	8.0	25.2	10.5	8.7
Spinach (<i>Beta vulgaris</i>)	Shoot	35.8	63.0	970	54.9	6.0	16.9	5.8	3.2
	Root	34.3	58.0	970	96.6	6.5	18.9	7.0	1.9
Jowar (<i>Sorghum bicolor</i>)	Shoot	23.3	140.3	644	51.0	5.0	22.2	4.7	0.9
	Root	39.8	128.9	1252	120.8	5.0	26.1	10.8	3.8
Coriander (<i>Coriandrum sativum</i> L.)	Shoot	35.8	66.0	552	18.7	3.5	17.3	3.7	0.4
	Root	12.8	85.5	967	66.3	5.0	22.4	6.5	3.0
Tomato (<i>Lycopersicon esculantum</i>)	Shoot	30.3	39.5	749	30.9	4.0	21.3	5.5	ND
	Root	22.8	107.6	1136	112.3	6.0	25.8	8.2	4.4
Brinjal (<i>Solanum melongena</i>)	Shoot	30.8	98.5	1049	76.0	5.5	68.9	7.2	3.9
	Root	15.3	103.5	1103	115.9	7.0	26.3	6.6	2.7
Longmelon (<i>Cucumis melo</i>)	Shoot	24.8	50.0	668	40.3	5.0	15.2	5.1	0.7
	Root	21.8	50.5	901	80.3	5.0	15.9	5.5	4.8

Table 1. Continued.

Karukshetra									
Chilli	Shoot	22.7	76.0	389	57.7	5.5	21.6	5.5	ND
<i>(Capsium annuum)</i>	Root	24.8	55.5	849	133.8	4.5	14.2	3.5	3.4
Kidney bean	Shoot	29.3	124.9	299	25.3	4.5	21.9	3.7	ND
<i>(Phaseolus vulgaris L)</i>	Root	32.8	69.0	1037	31.6	5.0	12.0	3.4	1.3
Tomato	Shoot	70.3	38.0	831	29.2	6.5	16.4	5.4	2.6
<i>(Lycopersiconesculantum)</i>	Root	29.3	199.9	1127	67.1	9.0	13.3	8.1	4.3
Onion	Shoot	15.3	137.5	879	52.3	4.5	19.9	4.9	ND
<i>(Alliumcopa)</i>	Root	35.8	113.2	925	88.7	6.0	60.9	6.2	2.5
Barseem	Shoot	16.3	103.8	647	42.8	5.5	12.7	5.0	0.8
<i>(Alexandriumtrifolium)</i>	Root	30.2	154.9	1284	288.7	7.0	39.1	12.9	10.4
Bitter gorud	Shoot	23.8	69.0	808	53.5	7.0	28.5	5.0	1.1
<i>(Momordicacharantia)</i>	Root	11.3	114.5	1094	169.6	18.0	32.3	7.8	4.0
Karnal									
Chilli	Shoot	28.8	62.0	1027	83.6	6.5	27.4	6.2	3.9
<i>(Capsium annuum)</i>	Root	26.3	103.5	1065	121.4	6.0	31.5	6.4	7.7
Barseem	Shoot	32.8	103.5	1040	89.9	6.0	21.6	8.1	2.2
<i>(Alexandriumtrifolium)</i>	Root	30.3	149.3	1266	229.8	7.0	39.8	12.8	8.2
Jowar	Shoot	34.8	38.0	867	34.5	4.0	6.5	3.6	3.5
<i>(Sorghum bicolor)</i>	Root	14.3	73.5	1227	91.6	5.0	15.2	7.7	2.9
Bottle gourd	Shoot	9.3	46.5	1052	55.1	5.5	23.3	7.7	2.8
<i>(Lagenoriasiceraria)</i>	Root	27.3	50.5	10093	110.5	4.5	15.1	6.8	6.2
Tomato	Shoot	28.3	42.0	85.7	54.2	6.0	18.8	6.9	3.8
<i>(Lycopersiconesculantum)</i>	Root	8.3	93.0	1055	69.4	6.0	26.6	6.7	4.1
Ridge gourd	Shoot	45.8	88.5	1078	96.6	6.0	19.0	8.8	2.8
<i>(Luffaacutangula)</i>	Root	37.8	136.2	1287	104.2	8.5	24.9	22.4	8.2
Panipat									
Water Melon	Shoot	11.8	45.0	238	41.4	2.5	9.4	3.4	ND
<i>(Citrullus vulgaris)</i>	Root	25.8	64.5	1084	116.3	6.5	13.6	8.7	3.3
Ridge Gourd	Shoot	19.5	65.5	462	32.6	5.5	12.0	1.9	0.6
<i>(Luffaacutangula)</i>	Root	13.5	46.5	879	129.0	7.0	29.0	7.7	4.5
Onion	Shoot	12.0	24.0	457	36.0	5.5	14.4	6.9	ND
<i>(Allium cepa)</i>	Root	7.5	32.5	606	59.8	6.0	18.0	5.5	0.8
Round Melon	Shoot	17.0	56.5	638	38.9	7.5	24.7	8.3	0.4
<i>(Citrullus vulgaris) var Fistulosus</i>	Root	18.5	96.5	640	87.5	5.5	17.0	6.3	0.9
Cauliflower	Shoot	7.0	49.0	306	43.6	6.0	16.3	4.7	ND
<i>(Brassica oleracea var. Botrytis)</i>	Root	13.5	59.5	555	40.0	6.0	14.3	5.1	2.2
Faridabad									
Okra	Shoot	1.2	6.0	90	24.8	3.0	4.9	1.8	ND
<i>(Ablemoschusesculantus)</i>	Root	14.0	37.0	464	73.1	6.5	16.2	7.6	0.5
Jowar	Shoot	14.0	39.5	427	76.0	6.0	19.6	5.7	ND
<i>(Sorghum bicolor)</i>	Root	17.5	99.0	1096	92.3	5.5	15.5	6.7	1.0
Brinjal	Shoot	25.0	53.0	719	49.6	6.5	17.4	5.6	0.5
<i>(Solanummelongena)</i>	Root	63.5	77.0	1039	79.8	6.5	22.6	9.7	2.8
Mint	Shoot	12.5	106.9	480	48.7	5.5	15.5	5.3	ND
<i>(Monthaarvensis)</i>	Root	15.0	104.5	966	42.7	6.0	14.8	9.5	3.2
Spinach	Shoot	16.5	46.0	681	106.9	6.5	17.1	9.6	0.3
<i>(Beta vulgaris)</i>	Root	26.0	50.5	786	136.0	6.5	14.6	9.0	1.6

REFERENCES

- Alawsy W.S.A., Alabad L.A.S., Hussein M.K. (2018) Effect of sewage water irrigation on growth performance, biomass and nutrient accumulation in *maize* and barley. *Int. J. Agric. Stat. Sci.* 14(2) : 519—524.
- Alghobar M.A., Suresha S. (2017) Evaluation of metal accumulation in soil and tomatoes irrigated with sewage water from Mysore city, Karnataka, India. *J. Saudi. Soc. Agric. Sci.* 16 : 49—59.
- Alloway W.H. (1968) Agronomic controls over the environmental cycling of trace elements. *Adv. Agron.* 20 : 235—274.
- Anwar S., Nawaz M.F., Gul S. (2016) Uptake and distribution of minerals and heavy metals in commonly grown leafy vegetable species irrigated with sewage water. *Environ. Monitor. Assess.* 6 : 188—541.
- Chaoua S., Boussaa S., Gharmali A., Boumezzough A. (2019) Impact of irrigation with wastewater on accumulation of heavy metals in soil and crops in the region of Marrakech in Morocco. *J. Saudi. Soc. Agric. Sci.* 18 : 429—436.
- Dadar M., Adel M., Ferrante M., Saravi H.N., Copat C., Conti G.O. (2016) Potential risk assessment of trace metals accumulation in food, water and edible tissue of rainbow trout farmed in Haraz River, Northern Iran. *Toxin. Rev.* 35 : 141—146.
- Khaled S., Muhammad B., Ashraf A. (2016) Field accumulation risks of heavy metals in soil and vegetable crop irrigated with sewage water in western region of Saudi Arabia. *Saudi. J. Bio. Sci.* 23 (1) : 532—544.
- Knezek B.D., Greinert H. (1971) Influence of soil Fe and Mn EDTA interaction upon the Fe and Mn nutrition of Bean plants. *Agro. J.* 63 : 617—619.
- Meena R., Datta S., Golui D. (2016) Long-term impact of sewage irrigation on soil properties and assessing risk in relation to transfer of metals to human food chain. *Environ. Sci. Poll. Res.* 23 : 14269—14283.
- Mortvedt J.J., Cox F.R., Shuman L.M., Welch R.M. (1991) Micronutrients in Agriculture. 2nd edn. Soil Sci. Soc. America, Inc., Madison, U.S.A.
- Naz S., Anjum M.A., Akhtar S., Naqvi S.A.H., Chohan S., Zulfiqar M.A. (2018) Effect of different irrigation sources on proximate composition and heavy metals uptake in some selected vegetables. *Pak. J. Agric. Res.* 31(4) : 336—346.
- Nazir R., Khan M., Masab M., Rehman H.U., Rauf N.U., Shahab S., Ameer N., Sajed M., Ullah M., Rafeeq M., Haheen Z. (2015) Accumulation of Heavy Metals (Ni, Cu, Cd, Cr, Pb, Zn, Fe) in the soil, water and plants and analysis of physico-chemical parameters of soil and water collected from Tanda Dam kohat. *J. Pharm. Sci.* 7 : 89—97.
- Rai S., Chopra A.K., Pathak C., Sharma D.K., Sharma R., Gupta P.M. (2011) Comparative study of some physico-chemical parameters of soil irrigated with sewage water and canal water of Dehradun city, India. *Appl. Sci. Res.* 3 : 318—325.
- Rajput S.G., Tiwari D., Katiyar D., Kumar N., Kumar A., Pathak R.K., Srivastava V. (2017) Effect of industrial effluent on soil properties, essential nutrient and pollutant element status of soils and plants in a vegetable growing near Jajmau industrial area of Kanpur, Uttar Pradesh, India. *Res. Crops.* 18 (2) : 249—255.
- Salakinkop S.R., Hunshal C. S. (2014) Domestic sewage irrigation on dynamics of nutrients and heavy metals in soil and wheat (*Triticum aestivum* L.) production. *Int. J. Rec. Org. Waste. Agric.* 3(64) : 1—11.
- Tandon H.L.S. (1995) Method of Analysis of soils, Plant, Waters and Fertilizers. Fertilizers Development and Consultation Organization, New Delhi.