Environment and Ecology 38 (3B) : 668—677, July—September 2020 ISSN 0970-0420

Plastic Waste Management in Asia Pacific Region : Issues and Challenges

Annes Elsa Francies, Sunil Herat

Received 7 July 2020; Accepted 24 August 2020; Published on 11 September 2020

ABSTRACT

Rapidly growing plastic usage and its subsequent release into the natural environment significantly increase the amounts of plastics entering oceans and require urgent action on a global scale. As the world's fastet-growing economics. Asia-Pacific countries have a very high material footprint as well as a plastic footprint. Massive urbanization, rapid industrialization, and changing production and consumption practices made many countries in the region as the world's most significant contributors of plastic litter. According to studies, around 60% of the total mismanaged plastic waste is coming from the region. In this paper, a few common challenges faced by these countries inadequately managing their plastic waste are identoified. This includes inefficient waste management systems, high rates of informal waste picking and treatment, and insufficient waste legislations. It is also found that the primary cause of marine plastic

Annes Elsa Francis

Sunil Herat

pollution is unsustainable human activities, and the significant pathways of land-based plastic litter to the oceans are rivers.

Keywords Marine Litter, Marine Plastics, Asia-Pacific Region, Plastic Waste Mangement, Plastic Leakage.

INTRODUCTION

Despite their high significance, oceans are now encountering various persisting and wide-reaching threats. Among this, the most pressing problem is marine litter arising from both land and sea-based souress. Marine debris consists of all objects that do not naturally present in the coastal environments such as the shorelines, seabed, water column, and water surface but are nevertheless found there (UN Environment Programme 2001). If not adequately managed, any human-made material that does not deteriorate naturally within a short period can become a source of marine litter (Grida. no 2016). Studies show that the most significant producers of marine debric are humans. Human activities like intentional or unintentional dumping of waste in freshwater sources and coastal environments and leaving unwanted materials in seashores and on beaches during recreational activities are considered as the principal causes of rising marine pollution. The abundance of consumer items like plastic food wrappers, plastic bags and bottles, straws, stirrers, cigarette butts, rubber thongs, and cosmetic products in the marine environments prove

Postgraduate Student, School of Engineering and Built Environment, Griffith University, Queensland 4111, Australia Email: anneselsa.francis@griffithuni.edu.au

Associate Professor, School of Engineering and Built Environment, Griffith University, Queensland 4111, Australia Email: s.herat@griffith.edu.au *Corresponding author

the massive involvement of humans in the generation and accumulation of marine litter (United Nations Sustainable Development 2016).

Plastics and plastic related materials constitute more than 80% of the total marine litter present in the environment today, and according to the UN, seven out of the top ten marine litter items are made of plastic polymers (United Nations Sustainable Development, 2016, Camey Almroth and fggert 2019).

It is expected that, by 2025, oceans might carry one tonne of plastic waste for every three tonnes of fish. Hence, if not carefull managed and no proper preventive measures are taken immediately, within five years, there will be more plastics in oceans than fish by weight, which will cause severe ecological, economic, cultural, and health impacts worldwide (Www3.weforum.org 2016). Besides, since the effects of plastic litter often occur in places that are far away from the points of its source, marine plastic pollution is becoming more sophistiated and unpredictable with each day.

Rapid population growth, that urbanization, quick industrialization, and changing lifestyles significantly increase the consumption of resources in the region, which in turn leads to enormous amounts of waste, mainly plastic waste. With the region's plastic waste production is expected to reach 140 million tonnes by 2030, Asia-Pacific is facing numberous challenges in managing their increasing amounts of plastic waste (UNCRD 2019). Most of the region's plastic waste except for developed countries like Australia, Japan, and Singapore, is not adequately managed. It is thus threatening to the land and coastal environments with linkages to livelohood problems. This mismanaged plastic is the primary source of global ocean plastic pollution, and its scale is a reason for serious concerns. More than 70% of the global mismanaged waste is coming from the Asia Pacific region, with the top eleven countries in which mismanage plastic waste is from the region. Besides, eight of the top ten litter contributing rivers are in Asia (Gray 2018, UNCRD0 2019). All of these factors make the region the center for marine plastic pollution. Hence, it is of fundamental importance to form systematized and well-coordinated strategies and measures to manage and prevent the production, consumption, and disposal of plastics in the asia Pacific countries. According to studies, by reducing 65% of plastic leakage from just five Asia Pacific countries: China, Indonesia, Philippines, Thailand and Vietnam can reduce fotal global plastic leakage by approximately 45% by 2025 (McKinsey & Company 2017, Shuker and cadman 2018).

This paper aims to assess the production, consumption, and management of plastic in the Asia Pacific region to find out the common issues faced by the region's plastic waste management sector by analyzing a few countries.

Plastic production and consumption

In 2019, annual global plastic production surpassed 380 million tons, almost the same as the weight of the entire human population (Plasticseurope.org 2019, Ritchie and Roser 2019). So far, a total of 8.3 billion tons of virgin plastics have been produced in the world using 4 to 8% of the world's total oil production, and among this, around 6300 metric tons came out as plastic waste since 2015. If the current trends and methods of production and waste management continue, the amount of plastic waste will reach almost 12,000 metric tons by 2050 and consumes 20% of the world's oil production (Geyer at al. 2017).

Two types of plastics run the plastic industry; Thermoplastics and Theemoseta; and the global plastic production is dominated by thermoplastics such as polypropylene (PP), low -and linear low-density polyethylene (LLDPE), high-density polyethylene (HDPE). polyvinyl chloride (PVC), polystyrene (PS) and polyethylene terephthalate (PET) mainly due to its reversible characteristics. These are also the major types of plastics used in the packaging industry, and therefore, they are prouced abundantly and consequently, making them the most likely ones to end up in the oceans (Andrady 2011, Hahladakis et al. 2018).

Only 14% of the total plastic packaging is recycled every year, and only 5% of their material value is kept for future use after sorting and reprocessing. This results in a loss of 80 to 120 billion USD each year to the economy. Besides, the rest of the packaging

Table 1.	Plastic	Waste in the	World's Oceans	(Gray 2018).

Ocean	Count (trillion)	Weight (tons)	
North Pacific	1.990	964.0	
South Pacific	0.491	210.2	
North Atlantic	0.930	564.7	
South Atlantic	0.297	127.8	
Indian Ocean	1.300	591.3	

materials can end up in our natural environments and oceans as pollutants (Ellen MacArthur Foundation 2017).

Only 14% of the total plastic packaging is recycled every year, and only 5% of their material value is kept for future use after sorting and reprocessing. This results in a loss of 80 to 120 billion USD each year to the economy. Besides, the rest of the packaging materials can end up in our natural environments and occeans as pollutants (ellen MacArthur Foundation 2017).

Marine plastic litter

Leakage of plastics into the natural environment can happen at any stage of its production-use-disposal cycle; however, it can reach oceans mainly through weak solid waste management systems and inadequate wastewater treatment processes (UNEP 2016). Three-fourths of the total plastic leakage comes from land-based sources and originates from uncollected rubbish, while the rest originates from within the waste collection systems itself (McKinsey and Company and Ocean Conservancy 2015). Table 1 shows the marine plastic litter distribution in world oceans.

As per the World Economic Forum, between 4.8 and 12.7 million metric tons of plastics reach oceans every year, making a current total of more than 150 million metric tons. This amount is equal to disposing of the rubbish of one garbage truck each minute into the ocean. Without proper action, this is expected to increase furthermore in the coming future as two per minute by 2030 and four per minute by 2050 (Ellen MacArthur Foundation 2017). In marine environments, plastics can undergo different types of degradation processes due to the combined action of atmospheric oxygen sunlight, and seawater.

These degradation processes can be of five kinds : Thermooxidative degradation, hydrolytic degradation, biodegradation, mechanical degradation, and photodegradation. among this, photodegradation is the most prevalent in coastal environments due to UV radiation's presence. However, degradation processes are alower within water due to a fall in temperature and limited UV penetration, which leads to the accumulation of plastic debris in ocean depths (Niaounakis 2017). However, the processes mentioned above disintegrate plastic waste into varying sizes, namely: macro, meso, micro, and nano plastics, and among these, the most common ones are microplastics. It can be of two types, primary and secondary. Primary incroplastics are those which are produced as microbeads, capsules, or fibers and commonly found in hygiene or cosmetic products or as plastic pellets that used as raw materials in the plastic industry or air-blasting media whereas the fragmentation of more abundant materials through the processes mentioned above produces secondary microplastics (Cole et al. 2011, Thompson 2016).

The size of microplasties are varying between studies, however many authors and the National Oceanic and Atmospheric Administration (NOAA) have defined it as particles smaller than 5 mm diameter (Barboza and Gimenez 2015, Van Cauwenberghe et al. 2013, Thompson 2016, Nguyen et al. 2019). This smaller size gives it a high surface area for sorption of contaminants and the potential to translocate in the bodies of organisms (Grida. no 2016, Nguyen et al. 2019). Its small sizemakes it challenging to identify and separate them from the environment, and hence, to understand its quantity, environmental prevalence, and potential threats became complicated. Therefore, plastics can remain in the oceans for hundreds or sometimes thousands of years in their original form and can persist even longer as disintegrated fragments and can go unnoticed (Grida. no 2016).

Impacts of plastic litter on marine animals and other aquatic life

Marine debris is not only sesthetically undesirable but also harmful to virtually all marine species and ecosystems. In a broader context, potential impacts of marine plastics can be categorized in three major

areas; injury or death to marine life, harm to the coastal and marine environment and dangerous to human health and economy (agamuthu et al. 2019) The first two are relevant in this paper and hence discussed in detail here. Impacts of plastic litter on marine life and the ecosystem are determined by type, size, quantity, and frequency of occurrence of debris (Rochman et al. 2016). however, these effects can be extremely dangerous and wide-reaching, and among them, the most noticeable, disturbing and harmful impact are related to ingestion and entanglement (Gall and Thompson 2015, Issues Brief - Marine Plastic 2018). Globally, more than 800 marine species, including seabirds, marine turtlies, whales, fishes and other reptiles are affected through the ingestion and entanglement of marine litter (Csiro.au 2019, United Nations Sustainable Development 2016). According to fata, the very first incident of entanglement was reported in Cojimar Bay in Habana Harvour in 1931, where a shark was seen entangled in a rubber automobile tire (Gudger 1931, van Truong and beiPing 2019). Since plastic and other synthetic materials materials became common in fishing equipment as well as in beach recreation activities, entanglement of marine animals in ALDFG, ghost fishing and single-use plastic products became a huge issue. Entanglement can restrict the marine animals which will lead to smothering, interruptions in feeding, starvation and eventually drowning. It can also result in physical trauma and infections in animals as the gear cut cut into their flesh, causing life-threatening injuries and ultimately death (NOAA 2014, NOAA Fisheries 2017).

Plastics that are highly persistent in marine conditions, particularly in the form of micro- and nano plastica can be ingested by marine life, especially seabirds and fishes, and can travel throughout the food chain. These plastic materials which contain many harmful chemicals like persistent organic pollutants andendocrine disruptors, can cause serious harm to marine animals (Gallo et al. 2018). accumulation of microplastic particles in marine animals can cause health issues such as blockages in the digestive system, starvation, and physical deterioration, which eventually results into a lack of reproductive fitness, weakness in feeding ability, reduced predator avoidance and even drowning (Wright et al. 2013). In addition to the internal accumulation of debris. external adsorption of plastic particles, specifically nanoparticles may obstruct algal photosynthesis by physically blocking air and sunlight while increasing reactive oxygen species production.Microplastics can lead to reduced availability of nutrients in microalgae (Bhattacharya et al. 2010, Prata et al. 2019).

Morcover, plastic debris can promore the spreading of nonindigenous species by supporting its growth by being a new safe habitat for them to survive. The distribution, characteristics and life span of plastic are different from that of the natural substrate like wood and microalgae, helping invasive species to develop, persist and even breed in them. Later these plastic debris items can act as a medium oftransport by taking flora and fauna attached to it into faraway places where it is foreign and can become aggressive and harm the local ecosystem (Garcia-Vazquez et al. 2018, GESAMP 2016.

Why it is important to tackle marine plastic litter

Covering almost 72% of the earth's surface, oceans regulate Earth's temperature and climate patterns, generate much of the oxygen we breathe, and provide food for more than three billion people in the world (UNEP, n.d.). While keepingpeople all around the globe connected and apart, oceans stay as the foundation for most of the world's largest economies (UNEP -UN Environment Programme 2019). However, the increasing amount of ocean litter and other unsustainable human activities taking place in marine environmenta prevent oceans from providing the above-said services. With more than 8 million tonnes of plastic reachin oceans every year, plastic litter cause irreversible damage to ocean life and marine ecosystems. Due to its durability, abundance and persistence, plastic litter has become a major environmental problem and a public health concem of our century. The volume of plastic coming into the sea are expected to increase in he coming years if countries around the world failed to take adequate measures to manage their growing rate of waste (Alpizar et al. 2020, Tibbetts, 2015). This will result in economic, cultural, aesthetic, environmental and health impacts worldwide, including degradation of marine and coastal habitats and ecosystems, which will causes substantial socioeconomic losses in maritime nations and marine industries (COBSEA 2018).

Marine plastics in asia Pacific region

According to studies, the top eleven countries which mismanage plastic waste comes from the Asia-Pacific region and are China, Indonesia. This Philippines, Vietnam, Sri Lanka, Thiland, Malaysia, Bangladesh, India, Pakistan and Myanmar. Mismanagement and resulting leakage of plastic waste into more extensive environments create immense pressure on the finite natural resources of the region, including aquatic and marine ecosystems. (UNCRD 2018, UNESCAP 2019). During 2014, Asia-Pacific's total annual municipal solid waste (MSW) generation was 870 million tonnes, with an average of 1.4 kg per person per day. This accounts for almost 43% of the total global MSW production, and within this plastic waste constitutes 8-12% (UNCRD 2019). The entire MSw generation of the region is expected to reach 1.4 nilliontonnes a year by 2030 with 1.6 kg per capita per day and the subsequent plastic waste generation is assumed to reach 140 million tonnes (UNCRd 2019) Within the region, East Asia and the Pacific countries are the biggest generators of waste in the world by producing 468 million tonnes of waste in 2016, at an average per capita rate of 0.56 kg per day. At the same time, South Asia generated 334 million tonnes of waste at an average of 0.52 kg per person per day. Almost half of the total global plastic is produced by Aia and is recognized as the world's largest waste-producing continent. Within Asia, China and Japan lead the plastic production with a rate of 29.4% and 3.95 of total plastic production. respectively, and the rest of Aiia produces 16.8% of global plastic (UNCRD 2019). It is estimated that 1.7 to 4.9% of plastic waste generated on land becomes marine litter. It this rate continues by 2030, 2.3 to 6.4 million tonnes of plastic waste will reach the oceans from the Aia-Pacific region alone (UNCRD 2019).

According to studies reducing 65% of leakage from five focus countries: China, Indonesia, Philippines, Thailand, and Vietnam can reduce total global plastic leakage by approximately 45% by 2025 (McKinsey & Company 2017, Shuker and Cadman 2018). Table 2 shows the contribution made to marine litter by top plastic generating countries in the Asia Pacific region.

Plastic waste management in selected countries

Indonesia

Plastic production in Indonesia is rising at an annual rate of 4% and has increased from 2.3 million tonnes in 2013 to 2.9 million tonnes in 2017. At the same time, yearly plastic consumption showed a growth rate of 6% and has risen from 4 million tonnes in 2013 to 4.9 million tonnes in 2017 (UNCRD 2019), 40% of Indonesia's total plastic demand is met by by importing plastic from foreign countries and the major plasic consuming sectors of the country are packaging, which assounts for 49%, followed by construction and automotive industries (UNCRD 2019).

Indonesia produces around 5.4 million metric tons of plastic waste every year at a rate of 0.52 kg per capita per day. Among this only 39% is collected on average, and in rural and remote areas, these rates

Country	Waste Generation Rate (kg/person/day)	Coastal Population (million)	Mismanaged waste (%)	Mismanaged Plastic waste (million tones/year)	Mismanaged Plastic Waste (%)	Plastic Marine debris (million tonnes/year)
Bangladesh	0.43	70.9	89	0.79	2.5	0.12-0.31
China	1.10	262.9	76	8.82	27.7	1.12-3.53
India	0.14	187.5	87	0.60	1.9	0.09-0.24
Indonesia	0.52	187.2	83	3.22	10.3	0.48-1.29
Malaysia	1.52	22.9	57	0.94	2.9	0.14-0.37
The Philippines	0.5	83.4	83	1.88	5.9	0.28-0.75
Vietnam	0.79	55.9	88	1.83	5.5	0.28-0.73
Thailand	1.2	26.0	75	1.03	3.2	0.15-0.417

Table 2. Estimated Quantities of Mismanaged Plastic Waste Entering the ocean from the Top Polluters (UNCRD 2018).

are as low as 16% (Pandjaitan 2020). Out of the total plastic waste generated every year, around 3.2 million metric tons are mismanaged, and approximately 1.25 million metric tonnes are estimated to reach oceans as marine litter making the country the second-largest donor of plastic waste into the oceans (Indonesia-investments 2019, McCarthy 2020, Syakti et al 2017). Composition of Indonesia's marine debris is found to be influenced by local lifestyles, and behavioral patterns and commonly contains synthetic materials like plastic bags, fishing accessories, food wrappers, bottles cans, footwear, polystyrene, polyethylene, and other remains of glass and plastics (Purba et al. 2019, Syakti et al. 2017, Plastic waste is not only foud in Indonesia's oceans, but also its rivers, and Indonesia is home to four of the top 20 polluting rivers in the world (Anne Jones 2019, Indonesia-invetments 2019, Purba et al. 2019).

China

As the world's largest manufacturing economy and exporter of goods, China is the largest producer of plastic with an average of 60 million tons per year, which accounts for nearly 29.4% of the total global plastics production (Ritchie and Roser 2018, Statista 2020, UNCRD 2019), Plastic production in China was growing at a rate of 7.4% annually and increased from 54.5 to 72.67 million tons between 2013 and 2017 while plastic consumption showed an annual growth rate of 5.1% and increased from 70.2 to 88.12 million tons during the period (Chen et al. 2020). In 2015, the nation's total plastic usage reached 1.4 times that of Europe and 2.6 times that of the United States. China recorded its highest material consumption of 35.2 billion metric tons in 2017 (Chen et al. 2020 UNCRD 2019).

China is also recognized as the world's largest trash generator by producing more than 15% of the world's total global municipal solid waste volume (Statista 2020, UNCRD 2019). Besides, the country is responsible for producing nearlt 30% of global plastic waste (Garcia et al. 2019). growing coastal population, increasing plastic usage, and inefficient and under-invested waste recycling sector are the main reasons behind China's growing plastic waste generation. Besides, once the largest importer of plastic waste, studies show that China had imported an aggregate of 45% of the global plastic waste between 1992-2016 (Garcia et al. 2019, Chen et al. 2020). Therapidly growing delivery and take away food industry in the country triggers a sudden rise in the nation's plastic usage. The packaging sector uses 42.1% of the total plastic resins. It is estimated that in 2017, the food industry consumed more than 60 million plasticcartons, and the food delivery processes used 14.7 billion plastic bags (chen 2018). The construction sector, with 22.4%, followed this, the automotive industry with 7%, the erectrical, electronics, and telecon sectors with 6.7% (UNCRD 2019)>

In China, rivers act as the significant pathways f plastic to oceans as regions near to estuaries with substantial river runoff and highly populated watersheds are major plastic litter hotspots. China's Yangtze River catchment is recognized as the most polluting river catchment in the world, and marine litter frequencies, rated as "extremely dirty" in the Yangtze Estuary, were almost 100% (Alkalay et al. 2007, Chen et al. 2020). Microplastics forms of polymers such as polyethylene, polyesters, polyethylene terephthalate, and polystyrene are the most abubdant types of plastics seen in China's marine waters (Qiu et al. 2015). According to the beach cleanliness assessment conducted by the Ministry of the Environment, Marine and Coastal Environment Division of Israel in 2018, almost half of China's beaches can be named as "extremely dirty beaches," and the ratio of it to "immaculate beaches" was only 8.5% (Alkalay et al. 2007).

Japan

Japan is the second-largest plastic producing country in Asia by providing 3.9% of total global plastic production (UNCRD 2019). In 2018, approximately 5.88 million tonnes of plastic products were produced in Japan. The major categories of processed plastics production are films and sheets, including bags, packaging, and sheeting for construction and building materials (statista 2019 abed, Yolin 2015).

Japanse total resin production in 2018 was 10,670-kilo tonnes, while domestic plastics consumption was 10.290-kilo tonnes. Complete plastic waste

created during the same year was 8.910-kilo tonnes, with 4,290-kilo tonnes of household plastic waste and 4620-kilo tonnes of industrial plastic wasste. The quantity of plastic waste that was effectively used was 7,5000 kilo tonnes making for an effective plastic utilization rate of 84% in the year (Plastic Waste Management Institute 2019). Japan has also been reported previously to export around 1,500 kilo tonnes of plastic waste per year to overseas destinations as a resource with the majority going to China (Plastic Waste Management Institute 2019). since the plastic import ban by China and other Asian countries, it has become a necessity to create more domestic resource recycling facilities, and this lead to many advancements in japan's plastic waste management system in the last few years (The sasakawa Peace Foundation 2019).

As an early adopter of the 3Rs (reduce-Reuse-Recycle), Japan handles most of its waste correctly. The primary recycling method used in Japan is mechanical recycling, which produces new plastic products from plastic waste (UNCRD 2019). The amount of plastic waste used in mechanical recycling as raw materials were 2.080-kilo tonnes in 2018, comprising of 710kilo tonnes (17%) of domestic plastic waste and 1370 kilo tonnes (30%) of industrial plastic waste (Plastic Waste Management Institute 2019).

Challenges facing Asia Pacific countries in managing plastic waste

When plastic waste is not effectively managed on land, it may enter natural waterways and can eventually reach oceans. Waste leakage hotspots are mainly found in three land-use types: Beach and recreation areas, human settlements; and industrial zones and urban areas. Settlements, particularly with less educated people and the residential regions with drainage channels connected to nearby rivers or canals, are most likely to pollute the water flow followed by illegal discharge from industries. These wastes are carried to the oceans mainly by waterways in the region. For example, the presence of waste leakage hotspots along riverbanks is familiarin countrieslike China and Indonesia. The Yangtze River of China takes an annual average of 0.33 million tonnes (33,000 tonnes) of plastic waste into the oceans. At the same time. Indonesia is home to four of the top 20 most polluting rivers in the world; Brantas, Solo, Serayu, and Progowhich respectively emit an average of 38,900, 32.500, 17,100 and 12,800 tones of plastic every year to occans (Anne Jones 2019, Indonesia-investments 2019, Purba et al. 2019).

Marine plastic litter can be identified as an essential part of the broader problem of poor waste management. When there is an absence of a proper waste management system in a region, waste reaches the oceans and causes pollution. Many Asia Pacific countries still lack adequate waste management systems in every stage, from the source of the waste to its final disposal. The significant challenges faced by the waste management systems of these countries are the lack of waste segregation and insufficient collection services. According to data, only 45 to 50% Indonesia's urban solid waste is collected effectively with significant variations among places, such as 98% collection and transfer in West Jakarta and 15% in South Tangerang. When it comes to plastics, out of the total 5.4 million metric tonnes of plastic waste produced annually, only 39% is collected, and in rural and remote areas, these rates are low as 16% (Pandjaitan 2020). Within the collected solid waste, around 81% goes uncorted, and another 10% is sorted but will end up remixed, making recycling processes difficult. Therefore, most of the plastic waste in the country, particularly from rural places, goes to landfills or leak into the ocean. Besides, Indonesia's waste management sector is heavily subsidized by local budgets. It receives only below-average funding of 5-6 dollars per capita per year allocations, where the international rates are 15-20 dollars per capita per annum. This limited funding is not enough for the proper strengthening of the waste management systems across the nation. Besides, this can cause a lot of inefficiencies, such as lack of infrastructure, failure to upgrade into modern techniques, and much higher operating costs for theavailable systems. Moreover, the lack of capacity in local governments significantly reduces confidence and increases risk among private investors.

Consumers play the most significant role in a country's waste management. Even though the population near coastlines as well as considerable litter

hotspots have diverse ethnicity, employmant status and duration of residency, communities from lower economic situations usually have poor and inadequate waste management services. Limited waste disposal facilities and services in these areas, together with a lack of knowledge and awareness about proper waste disposal methods among the residents, increase the magnitude of the problem.

Many studies suggest thatpeople with sducation and with higher environmental attitudes show more robust plastic avoidance. Moreover, the majority of mismanaged plastic comes from househol waste and food packaging, including bottles, bags, and eating accessories. Therefore, poviding consistent public information or campaigning about waste management can help in tackling plastic waste leakage (Heidbreder et al. 2019).

Vast quantities of plastic in waterways and oceans indicates that current systems of waste govermance are not good enough for the effective management of plastic waste. This failure of governance includes a lack of appropriate legal and regulatory frameworks for waste management, insufficient taxation systems for waste disposal, and failed enforcement of available laws and policies. Besides, current strategies and plans show no or very little importance to the waste issues in the upstream sides of pollution, such as river basins, which is the crtical pathway for plastic leakage to the oceans. Moreover, current regulations, policies, investments, and other waste management measures are distributed among various sectors and stakeholders and lack coherence.

Both formal and informal sectors performsolid waste collection in Aiia Pacific countries; however, the informal collectionconstitutes the majority. These waste pickers form the foundation of the informal waste management sector in these countries. Unfortunately, the unmanaged waste collection activities performed by people with less knowledge about the consequences of unsafe waste management is a challenge to implement a standardized waste treatment system in these countries. Waste pickers usually sort out the collected trash and left the unwanted materials on landfills, open dumpsites, or sometimes even in the streets, which can leak ito drainage systems and other waterways and can reach oceans in the end. Besides, rummaging through the waste without proper safety measures can cause severe health issues in these waste collectors (Zhang et al. 2010).

CONCLUSION

Poor and inadequate waste management practices and the failure to appreciate the potential uses of plastic waste as raw materials for new products along with the lack of awareness among the public regarding the consequences casually discarding plastics in natural environments fuelled the growth of marine plastic pollution in the past few decades. Since plastics are made in various diversities and compositions to last forever, it is more likely to rmain as a significant environmental problem for the coming decades. Today, the dangers of ocean plastics are restricted to marine life, or marine ecosystems and the potential threats of plastics are finding its way back to humans through the food chain.

Preventing plastic from entering the natural environments is likely to be far less costly and practical than getting rid of it once they are there. The first step to achieve this is to avoid the overall plastic contamination by reducing the production and use of plastic products followed by reusing and recycling plastic materials as long as possible and developing efficient waste management systems. This requiressignificant changes in the current production and consumption practices of plastics as well as need enhanced collaboration among different countries, organization, businesses, and individuals.

REFERENCES

- Agamuthu P, Mwhran S, Norkhairah A, Norkhariyah A (2019) Marine debris : A review of impacts and global initiatives. Waste Manag Res 37 (10) : 987—1002.
- Alakalay R, Pasternak G, Zask A (2007) Clean-coast index— A new approach for beach cleanliness assessment. Ocean and Coastal Manag 50 (5-6): 352—362.
- Alpizar F, Carlsson F, Lanza G, Carney B, Daniels R, Jaime M, Ho T, Nie Z, Salazar C, Tibesigwa B, Wahdera S (2020) A framework for selecting and designing policies to reduce marine plastic pollution in developing countries. Environm Sci & Policy 109: 25—35.
- Andrady A (2011) Microplastics in the marine environment. Marine Pollut Bull 62 (8) : 1505—1605.
- Anne Jones C (2019) Indonesia's Journey to Reduce 70% of Marine

Waste by 2025. [online] Asian Studies Center. Available at: https://asc,fisipol.ugm.ac.id/2019/06/18/indonesias-journey-to-reduce-70-of-marine-waste-by-2025/ [Accessed 14 Nov. 2019].

- Barboza L, Gimenez B (2015) Microplastics in the marine environment: Current trends and future perspectives. Marine Pollut Bull 97 (1-2) 5—12.
- Bhattacharya P, Lin S, Turner J, Ke P (2010) Physical adsorption of charged plastic nanoparticles affects algal photosynthesis. The J Physical Chem C 114 (39) : 16556—1656.
- Carney Almroth B, Eggert H (2019) Marine plastic pollution Sources, impacts, and policy issues. Rev Environm Econ Policy 13 (2): 317—326.
- Chen A (2018) China generates more waste than any other country. How does it deal with it ?, [online] World Economic Forum. Available at :https://www.weforum.org/agenda/2018/12/ no-chopsticks-with-may-takeaway-how-china-is-tacklingfood-waste-with-digital-innovation/ [Accessed 14 dec. 2019].
- Chen H, Wang S, Guo H, Lin H, Zhang Y (2020) A nationwide assessment of litter on China's beaches using citized science data. Environm Pollut 258 : 113756.
- Cole M, Lindeque P, Halsband C, Galloway T (2011) Microplastics as contaminants in the marine enbironment : A review. Marine Pollut Bull 62 (12) : 2588—2597..
- CSIRO (2019) Marine debris Sources, distribution and fate of plastic and other refuse – and its impact on ocean and coaastal wildlife. [online] Available at : Page 75 of 85. https://www.csiro.au/–/media/OnA/Files/15-
- Ellen MacArthur Foundation (2017) The New Plastics Economy: Catalysing Action. [online] Davos : World Economic Forum and Ellen MacArthur Foundation. Available at : https://www.ellenmacarthurfoundation.org/publications [Accessed 6 March 2020].
- Call S, Thompson R (2015) The impact of debris on marine life. Marine Pollut Bull 92 (1-2) : 170—179.
- Gallo F, Fossi C, Weber R, Santillo D, Sousa J, Ingram I, Nadal A, Romana D (2018) Marine litter plastics and microplastics and their toxic chemicals components: The need for urgent preventive measures. Environm Sci Europe 30 (1) In press.
- Garcia B, Fang M, Lin J (2019) Marine plastic pollution in Asia: All Hands on Deck !. Chinese J Environm Law 3 (1) 11–46.
- Garcia-Vazquez E, Cani A, Diem A, Ferreira C, Geldhof R, Marquez L, Molloy E, Perché S (2018) leave no traces – Beached marine litter shelters both invasive and native species. Marine Pollut Bull 131 : 314—322.
- GESAMP (2016)"Sources, fate and effects of microplastics in the marine environment: Part two of a global assessment" (Kershaw, P. J., and R\$ochman, C. M., eds).
- National Oceanic and Atmospheric Administration Marine Debris Program (2014) Report on the Entanglement of Marine Species in Marine Debris with an emphasis on Species in the United States. Silver Spring, MD, pp 28.
- Niaounakis M (2017) Degradation of Plastics in the Marine Environment. Management of Marine Plastic Debris. pp 127—142.
- Nguyen B, Claveau-Mallet D, Hemandez L, Xu E, Farner J, Tufenkji N (2019) Separation and analysis of microplastic and nanoplastics in complex environmental samples. Accounts of Chem Res 52 (4) : 858–866.
- NOAA Fisheries (2017) Entanglement of Marine Life : Risks and

Response. [online] Available at; https://www.fisheries.noaa. gov/insight/entanglement-marine-life-risksand-response [Accessed 22June 2020].

- Pandjaitan L (2020) Here's how Indonesia plans to tackle its plastic pollution problem. [online] World Economic Forum. Available at :https://www.weform.org/agenda/2020/01/heres-how-indonesia-plans-to-tackle-its-plastic-pollution-chal lenge/ [accessed 15 Fev 2020].
- Purba N, Handyman D, Pribadi T, Syakti A, Pranowo W, Harvey A, Issan Y (2019) Marine debris in Indonesia : A review of research and status.Marine Pollut Bull 146 : 134—144.
- Plasticseurope.org. (2019) Plastics the Facts 2019. [online] Available at : https://www.plasticseurope.org/application/ files/1115/72236/4388/FINA_web_version_Plastics_the_ facts2019_14102019.pdf[Accessed 5 Dec. 2019].
- Plastic Waste Management Institure (2019) An Introduction to Plastic Recycling 2019. [online]. Available at : http://www. pwmi.or.jp [Accessed 16 Dec. 2019].
- Prata J, da Costa J, lopes I, Duarte A, Rocha-Santos T (2019) Effects of microplastics on microalgae populations : A critical review. Sci of The Total Environ 665 : 400–405.
- Qiu Q, Peng J, Yu X, Chen F, Wang J, Dong F (2015) Occurrence of microplastics in the coastal marine environment : First observationon sediment of China. Marine Pollut Bull 98 (1-2): 274—280.
- Ritchie H, Roser M (2019) Plastic Pollution. [online] Our World in Data. Available at : https:/ourworldindata.org/plastic-pollution#share-of-global-total-mismanaged-plastic-waste-bycountry [Accessed 1 Dec. 2019].
- Rochman C, Prowne M, Underwood A, van Franeker J, Thompson R, AmaralZettler L (2016) The ecological impacts of marine debris: Unraveling the demonstrated evidence from what is perceived. Ecology 97 (2): 302–312.
- Shuker I, Cadman C (2018) Indonesia Marine debris hotspot rapid assessment: synthesis report. World Bank. Available at: http://documents.workldbank.org/curated/en/ 983771527663689822/Indonesia-Marine-debris-hotspot-rapid-assessment-synthesis-report.
- Statista (2019a) China: Amount of disposed waste 2018 / Statista. [online] Available at : https://www.statista.com/statistics/ 279117/amount-of-disposed-garbage-in-china/ [accessed 8 Jan. 2020].
- Statista (2020) China: plastic products production by month 2019 / statista. [online] Available at : https://www.statista. com/statistics/226239/production-of-plastic-products-in-chin-by-month/ [Accessed 5 Feb. 2020].
- Statista (2020b) Global plastic production /Statista. [online] Available at : https://www.statista.com/statistics/282732/ global -production-of-plasties-since-1950/[Accessed 4 Dec. 2019].
- Statista (2019c) Plastics products Japan by category 2018. [online] Available at: https://www.statista.com/statistics/486152/ processed-plastice-production-by-category-in-japan/[Accessed 2 Jan. 2019].
- Statista (2019d) Japan: plastic products production volume 2018 [Statista. [online] Available at: https://www.statista.com/ statistics/858647/japan-plastic-products-productionvolume/ Accessed 2 Jan. 2020].
- Syakti A, Bouhroum R, Hidayati N, Koenawan C, Boulkamh A, Sulistyo I, Lebarillier S, Akhlus S, doumenq P, Wong-Wah-

Chung P (2017) Beach macro-litter monitoring and floating microplastic in a coastal area of Indonesia. Marine Pollut Bull 122 (1-2) : 217—225.

- Thompson R, Olsen Y, Mitchell R, Davis A, Rowland S, John A, Mcgonigle D, Russell A (2004) Lost at Sea : Where IS All the Plastic? Science 304 (5672) : 838—838.
- Tibbetts J (2015) Managing Marine Plastic Pollution : Policy Initiatives to Address Wayward waste. Environm Hlth Perspectives 123 (4) In press.
- UNCRD (2019) Reducing plastic wastes through circular economy – Implications towards SDG 12 & SDG 14. [online} Available at :http:www.uncrd.or.jp/contet/documents/7608 PS-2-PPT-1-Background%20paper_Combined.pdf [Acccessed 8 Jan.2020].
- UUCRD (2018) State of the 3Rs in Asia and the Pacific -Experts' Assessment of progress in Ha Noi 3R Goals [online] Abailable at: http://www.uncrd.or.ip/state_of-3RS [Accessed 8 Nov. 2019].
- UN Environment Programme (2001) Marine Litter Trash That Kills. [online] Available at : https://wedocs.uncp.org/bitstream/handle/20.500.11822/9691/-Marine%20Liter%20-%20trash%20that%20kills-2003707.pdf?/sequence=3&%3BisAllowed=[Accessed 5 Nov. 2019].
- Unervironment,org (2018) Marine Litter and Plastic Pollution / Coordinating Body On The Seas Of East Asia (COBSEA). [online] Available at : https://www.unenvironment.org/cobsea/what-we-do/marine-litter-and-plastic-pollution [Accessed 22 June 2020].

UNEP (n.d.) GOAL 14: Life below water. [online] Available at :

https://www.unenvironment.org/explore-topics/sustainavledevelopmentgoals/why-do-sustainable-development-goalsmatter/goal-14.

- UNESCAP (2019) Closing the Loop Regional Policy Guide. [online] Available at : https://www.unescap.org/sites/default/ files/closing%20The%20Loop_Regional%20P olicy%20 Guide.pdf [accessed 10 Feb. 2020].
- Van Cauwenberghe L, Vanreusel A, Mees J, Janssen C (2013) Microplastic pollution in deep-sea sediments. Environm Pollut 182 : 495—499.
- van Truong N, beiPing C (2019) Plastic marine dobris ; Sources, impacts and management. Int J Environm Studies 76 (6) : 953—973.
- Wright S, Thompson R, Galloway T (2013) The physical impacts of microplastics on marine organisms : A review. Environm Pollut 178 : 483—492.
- Www3.weforum.org. (2016). The New Plastics Economy Rethinking the future of plastics. [online] Available at : http://www3.weforum.org/docs/WEF_The_Plastics_Economy.pdf [accessed 26 Nov. 2019].
- Yolin C (2015) Waste Management and Recycling in Japan Opportunities for European Companies (SMEs focus) / EU-Japan. [online] EU-Japan Centre. Available at : https://www. cu-japan.cu/publications/waste-management-and-recyclingjapan-opportunities-european-companies-smes-focus [Accessed 8 Oct. 2019].
- Zhang D, Tan S, Gersberg R (2020) Municipal soil waste management in China : Status, problems and challenges. J Environm Manag 91 (8) : 1623—1633.