

## **Growth and Productivity of Wicker Willow (*Salix triandra* L.) Plantation in Kashmir**

TAREQ AHMAD RATHER, K. N. QAISER AND M. A. KHAN

*Division of Forestry, Sher-e-Kashmir University of Agricultural Sciences &  
 Technology of Kashmir  
 Shalimar, Srinagar 191121, Jammu and Kashmir, India  
 E-mail : navedmoon2@hotmail.com*

### **Abstract**

An experiment on growth and productivity of wicker willow plantation (*Salix triandra* L.) was carried out on farmers field at Batipora (Taibal) in Srinagar district of Kashmir in India. The experiment was laid out in randomized block design with four treatments—fresh plantation ( $T_1$ ), two year old coppice ( $T_2$ ), three year old coppice ( $T_3$ ), four year old coppice ( $T_4$ ), each replicated six times. The initial nutrient status of the soil showed pH neutral to acidic, EC medium, organic carbon high with available NPK, low, high, medium, respectively. Planting distance of 15 × 30 cm was maintained in all the treatments. Sprouting vigor was found maximum in  $T_3$  (three year old coppice) and  $T_4$  (four year old coppice) which started and completed earlier than  $T_2$  (two year old coppice) and  $T_1$  (fresh plantation). Maximum growth and productivity was observed with different age of coppice (i.e. two year and four year old coppice, respectively). The number of shoots per coppice was the deciding factor in contributing yield, number of shoots (7.66) and dry yield (25.10 t/ha). In post-harvest processing rods were boiled in water for 10—12 hours and cooled for another 10—12 hours and peeling was done, the bark contributed nearly 25% of weight. The dimensions of processed rods falls into medium category in length and thick category in diameter based on the standards.

**Key words :** Growth, Productivity, Processing, Grading, *Salix triandra* L.

Willows are the fast growing pioneer species, their allocation of the assimilate transport is concentrated to the growing points of the shoots in the early stages of life (juvenile phase) and later willows invest more and more in the root system. This phenomenon is one of the reasons for early harvesting of willows to maintain the juvenile stages. As a result there is high woody biomass production in the stems of willows (1). Willow have been used by people for many centuries to build shelters, make fences, boats, agricultural implements, tooth picks, preparation of artificial limbs and baskets (2). Willows are easily propagated by stem cuttings. The cuttings are prepared out of vigorous 2—3 years old shoots free from disease and defects. These are obtained when the trees enter in the dormancy stage and are cut to the size of 20 cm length and mid point diameter 10—20 mm. Cuttings obtained in February to March and prepared from center half of the shoot.

Forest officers of Kashmir imported famous English bat willow (*Salix alba* cv Calva=*Salix coerulea*,

Smith) from England in 1917 and raised it in Kitreteng wetland nursery (Bijbehara Forest Division of Kashmir). They also imported some varieties of wicker willows from England such as *Salix triandra*, *Salix purpurea*, *Salix amygdalina*, *Salix hypopofolia*, *Salix viminalis*, *Salix flabellaries*. These are not planted by the departmental agencies as these plantations can be found in Harn, Shalbugh and other areas of Tehsil Ganderbal in the private lands of farmers raised for wicker works where water is readily available (2).

The *Salix triandra* L. (Almond willow), most commonly planted wicker willow species in Kashmir is a deciduous shrub upto 5 m height rarely a small tree, bark reddish flaking off in autumn like that of Chinar, leaves 5—10 cm, narrow, finely toothed, dark green and shining above, bluish or green beneath. The leaves have a distinct smell and taste of sweet almonds (hence known as almond willow). Winter twigs are brownish green shining above, buds narrow pointed, flowers March—May and some times

**Table 1.** Initial nutrient status of the soil.

| Nutrients  | Depth 0—30 cm                           |  |  |  | Range                     | Method employed       |
|--|---|--|--|--|---------------------------|-----------------------|
|  | T <sub>1</sub><br>(fresh<br>plantation) | T <sub>2</sub><br>(2 yr<br>old<br>coppice) | T <sub>3</sub><br>(3 yr<br>old<br>coppice) | T <sub>4</sub><br>(4 yr<br>old<br>coppice) |                           |                       |
| pH (1 : 2.5)                                       | 6.56                                    | 6.25                                       | 6.10                                       | 6.03                                       | Neutral to<br>mild acidic | Jackson (6)           |
| EC ds/m  | 0.16                                    | 0.18                                       | 0.22                                       | 0.23                                       | Medium                    | Jackson (6)           |
| OC (%)   | 1.14                                    | 1.40                                       | 1.65                                       | 1.78                                       | Medium to high            | Walkley and Black (7) |
| Available N (kg/ha)                                | 236.50                                  | 213.50                                     | 217.50                                     | 231.00                                     | Low                       | Subbiah and Asija (8) |
| Available P (kg/ha P <sub>2</sub> O <sub>5</sub> ) | 28.66                                   | 33.00                                      | 36.16                                      | 36.16                                      | High                      | Olsen et al. (9)      |
| Available K (kg/ha K <sub>2</sub> O)               | 224.00                                  | 223.66                                     | 224.50                                     | 225.33                                     | Medium                    | Jackson (6)           |

July and August. Catkins 2.5 to 5 cm long, male catkin with three stamens (3). The long branches of the almond willow are cut out after one year growth and processed in boiling water, then their bark is peeled off to make the rods water proof and more durable and the wickers are provided as a raw material to small scale industries of Kashmir for the manufacturing of chairs, tables, sofasets, baskets of various shapes and designs, flower vases and a variety of other fancy items (Fig. 1) (2).

Due to its fast growth, easy rooting, recurrent harvest and having commercial value for wicker works in Kashmir valley, the wicker willow *Salix triandra* L. is preferred over the main food crop of the valley i.e. paddy, due to lack of care and high returns. Till date no study has been done on wicker willow cultivation in Kashmir. Therefore, the present study was undertaken on the growth and productivity of wicker willow plantation in Srinagar district of Kashmir, where wicker willow is cultivated on wet sites. The survey of the area revealed that wicker willow has been found in all the districts of Kashmir except Kupwara. Also Srinagar district was found in the fore front of wicker willow cultivation (maximum families, area, yield and income).

(We acknowledge the chance for undergoing PG program by the Vice Chancellor, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Special thanks are due to Khawaja Naved Sahib and Manzoor Ahmad Khan for teaching assistance in field work and chemical analysis).

### Methods

The experiment was carried out on the farmers

field during 2005 at Batipora (Hazratbal), which is located at an altitude of 1,587 m asl. with a latitude of 34.08 °N and 74.83 °E. During the growing period the maximum mean temperature was recorded in June (30.72 C) and minimum in November (−0.80 C), rainfall was highest in July (28.3 mm) and minimum in November (3.05 mm) and relative humidity was maximum in October (78.02%) and minimum in June (58.56%).

The data on different parameters for growth and productivity of *Salix triandra* L. commonly known as wicker willow were collected from four treatments T<sub>1</sub> (fresh plantation) which was planted in the first week of March and remaining treatments—two year old coppice (T<sub>2</sub>), three year old coppice (T<sub>3</sub>) and four year old coppice (T<sub>4</sub>) were earlier harvested, once, twice and thrice. The common planting distance was maintained at 30×15 cm. Each treatment was replicated six times in randomized block design.

The number of coppice counted from the marked plots (1 m<sup>2</sup>) and were termed as plant density/m<sup>2</sup>. The coppice in marked sample plot was tagged and numbered one, two, three, the sample plots were visited

**Table 2.** Sprouting percentage of wicker willow (*Salix triandra* L.).

| Treatments                        | Per cent sprouting |           |           |           |          |
|-----------------------------------|--------------------|-----------|-----------|-----------|----------|
|                                   | 15<br>Mar          | 20<br>Mar | 25<br>Mar | 30<br>Mar | 4<br>Apr |
| T <sub>1</sub> (fresh plantation) | 1                  | 25        | 75        | 98        | 100      |
| T <sub>2</sub> (2 yr old coppice) | 2                  | 40        | 84        | 99        | 100      |
| T <sub>3</sub> (3 yr old coppice) | 2                  | 42        | 88        | 100       | 100      |
| T <sub>4</sub> (4 yr old coppice) | 3                  | 44        | 90        | 100       | 100      |
| Mean                              | 2                  | 37.75     | 84.25     | 99.25     | 100      |
| CD (0.05)                         | 0.12               | 4.06      | 6.35      | NS        | NS       |

**Table 3.** Average number of shoots/coppice of wicker willow (*Salix triandra* L.) and mortality (%).

| Treatments                        | Average number of shoots |        | Mortality (%) |
|-----------------------------------|--------------------------|--------|---------------|
|                                   | 15 Apr                   | 15 Nov |               |
| T <sub>1</sub> (fresh plantation) | 1.36                     | 1.30   | 4.41          |
| T <sub>2</sub> (2 yr old coppice) | 3.00                     | 2.86   | 4.66          |
| T <sub>3</sub> (3 yr old coppice) | 4.57                     | 4.31   | 5.68          |
| T <sub>4</sub> (4 yr old coppice) | 7.66                     | 6.97   | 9.00          |
| CD (0.05)                         | 0.33                     | 0.30   | 0.95          |

after five-day intervals to record the sprouting percentage. The shoots were counted from the randomly tagged coppice and were expressed as shoots/coppice. The shoot height and collar diameter were recorded at an interval of one month from the date of shoot emergence in cm from the randomly tagged coppice stands of 1 m<sup>2</sup> with the help of metallic scale vernier callipers. Also the number of leaves/shoots was recorded at an interval of one month in 1 m<sup>2</sup> sampling plot from the randomly tagged coppice and were expressed as leaves/shoot.

The maturity indices as perceived by the farmers was taken as harvesting stage and period of harvesting as harvesting time. From each sampling plot shoots were taken from an area of 0.25<sup>2</sup> m to determine the fresh and dry weight of shoots prior to leaf fall and after leaf fall and expressed as kg/0.25 m<sup>2</sup> which was converted to fresh weight (tonnes/ha). The shoots taken were oven dried at a temperature 105 ± 2 C for 72 hours and weighed (4). The weight obtained was reported as dry weight kg per 0.25 m<sup>2</sup> and then converted to dry weight t/ha. The moisture percentage in shoots prior and after leaf fall was determined by the formula :

$$\text{Percent moisture} = \frac{\text{Fresh weight} - \text{dry weight}}{\text{Fresh weight}} \times 100$$

After harvest samples were boiled in standard sized boiling tanks (4.57 m × 1.52 m × 1.52 m) made of cast iron and heated directly by fuel wood for 10–12 hours and then left for cooling for another 10 to 12 hours. After cooling samples were peeled by fastening the bark of rods between two wooden sticks held at cross with each other and partly embedded in the soil. Then the rods were air dried under sunlight for a

**Table 4.** Average shoot height of wicker willow (*Salix triandra* L.).

| Treatments        | Average shoot height (cm)         |                                   |                                   |                                   | Monthly increment (cm) |       |
|-------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|------------------------|-------|
|                   | T <sub>1</sub> (fresh plantation) | T <sub>2</sub> (2 yr old coppice) | T <sub>3</sub> (3 yr old coppice) | T <sub>4</sub> (4 yr old coppice) |                        |       |
| 15 Apr            | 19.28                             | 20.96                             | 20.52                             | 19.83                             | 20.15                  | –     |
| 15 May            | 47.04                             | 60.80                             | 54.01                             | 53.79                             | 53.91                  | 33.76 |
| 15 Jun            | 84.12                             | 107.05                            | 95.49                             | 88.53                             | 93.80                  | 39.89 |
| 15 Jul            | 113.13                            | 167.84                            | 155.51                            | 141.52                            | 149.50                 | 55.7  |
| 15 Aug            | 174.42                            | 188.30                            | 178.92                            | 160.76                            | 175.60                 | 26.1  |
| 15 Sept           | 189.21                            | 195.12                            | 183.96                            | 165.80                            | 183.52                 | 7.92  |
| 15 Oct            | 194.29                            | 197.87                            | 186.58                            | 170.62                            | 187.34                 | 3.82  |
| 15 Nov            | 194.58                            | 198.06                            | 186.64                            | 171.27                            | 187.64                 | 0.30  |
| Mean              | 129.51                            | 142.00                            | 132.71                            | 121.52                            | –                      | –     |
| CD (0.05)         | Month = 5.75                      |                                   |                                   |                                   |                        |       |
| Treatment         | = 4.07                            |                                   |                                   |                                   |                        |       |
| Month × treatment | = 11.51                           |                                   |                                   |                                   |                        |       |

weeks time, which was then graded depending upon the diameter and height in cm based on the standards into thin, medium and thick (5) with metallic scale and vernier calliper.

The soil samples were taken randomly from all the coppice stands of different ages at 0–30 cm soil depth. The samples were air dried under shade, crushed and passed through 2 mm sieve. The sieved samples were analyzed for different chemical properties. The methods used for analysis of various chemical properties are as follows :

| Chemical properties          | Methods employed      |
|------------------------------|-----------------------|
| pH (1 : 2..5)                | Jackson (6)           |
| EC (ds/m)                    | Jackson (6)           |
| Organic carbon (%)           | Walkey and Black (7)  |
| Available nitrogen (kg/ha)   | Subbiah and Asija (8) |
| Available phosphorus (kg/ha) | Olsen et al. (9)      |
| Available potassium (kg/ha)  | Jackson (6)           |

## Results and Discussion

The studies conducted on the soil analysis revealed that the soil pH decreased and organic carbon increased with the increase in age of the coppice (Table 1). The pH recorded was suitable for the growth of wicker willow (*Salix triandra* L.). The increase in organic carbon and decrease in pH was due to the reason that leaves of the deciduous trees decompose readily, which increased the organic carbon and dur-

**Table 5.** Average collar diameter of wicker willow (*Salix triandra* L.).

| Treat-<br>ments      | Average collar diameter (cm)                 |  |  |  | Monthly<br>in-<br>cre-<br>ment<br>(cm) |      |
|----------------------|--|--|--|--|--|------|
|                      | T <sub>1</sub><br>(fresh<br>planta-<br>tion) | T <sub>2</sub><br>(2 yr<br>old<br>coppice) | T <sub>3</sub><br>(3 yr<br>old<br>coppice) | T <sub>4</sub><br>(4 yr<br>old<br>coppice) |  | Mean |
| 15 Apr               | 0.09   | 0.11                                       | 0.10                                       | 0.10                                       | 0.10                                   | –    |
| 15 May               | 0.23   | 0.30                                       | 0.26                                       | 0.26                                       | 0.26                                   | 0.16 |
| 15 Jun               | 0.50   | 0.78                                       | 0.64                                       | 0.60                                       | 0.63                                   | 0.37 |
| 15 Jul               | 1.17   | 1.33                                       | 1.27                                       | 1.20                                       | 1.24                                   | 0.61 |
| 15 Aug               | 1.44   | 1.52                                       | 1.39                                       | 1.30                                       | 1.41                                   | 0.17 |
| 15 Sep               | 1.55   | 1.64                                       | 1.42                                       | 1.33                                       | 1.48                                   | 0.07 |
| 15 Oct               | 1.62   | 1.65                                       | 1.43                                       | 1.35                                       | 1.51                                   | 0.03 |
| 15 Nov               | 1.63   | 1.65                                       | 1.43                                       | 1.35                                       | 1.51                                   | 0.00 |
| Mean                 | 1.03   | 1.12                                       | 0.99                                       | 0.94                                       | –                                      | –    |
| CD (0.05)            |  |  |  |  |  |      |
| Month=0.10           |  |  |  |  |  |      |
| Treatment=0.07       |  |  |  |  |  |      |
| Month × treatment=NS |  |  |  |  |  |      |

ing decomposition of leaves the acids were released which consequently decreased the soil pH. These results were in agreement with earlier reports (10–12). The electrical conductivity increased with the increase in age of the coppice and was found in the range of medium to high (Table 1). EC was minimum (0.16 ds/m) in T<sub>1</sub> (fresh plantation) and maximum in T<sub>4</sub> (four year old coppice) (0.23 ds/m) which was best for the growth and development of the willows which are in conformation with earlier results (13).

**Table 6.** Average number of leaves/shoot of wicker willow (*Salix triandra* L.).

| Treat-<br>ments        | Average number of leaves                     |  |  |  | Monthly<br>in-<br>cre-<br>ment<br>(No) |       |
|------------------------|--|--|--|--|--|-------|
|                        | T <sub>1</sub><br>(fresh<br>planta-<br>tion) | T <sub>2</sub><br>(2 yr<br>old<br>coppice) | T <sub>3</sub><br>(3 yr<br>old<br>coppice) | T <sub>4</sub><br>(4 yr<br>old<br>coppice) |  | Mean  |
| 15 Apr                 | 7.71   | 8.38                                       | 8.23                                       | 7.95                                       | 8.07                                   | –     |
| 15 May                 | 1.81   | 24.31                                      | 21.60                                      | 21.51                                      | 21.56                                  | 13.49 |
| 15 Jun                 | 33.64  | 42.81                                      | 38.19                                      | 35.40                                      | 37.51                                  | 15.95 |
| 15 Jul                 | 53.91  | 67.13                                      | 62.20                                      | 56.60                                      | 59.96                                  | 22.45 |
| 15 Aug                 | 69.76  | 75.31                                      | 71.56                                      | 64.30                                      | 70.23                                  | 10.27 |
| 15 Sep                 | 75.67  | 78.04                                      | 73.58                                      | 66.31                                      | 73.40                                  | 3.17  |
| 15 Oct                 | 77.71  | 79.14                                      | 74.63                                      | 68.24                                      | 74.93                                  | 1.53  |
| 15 Nov                 | 77.82  | 79.22                                      | 74.65                                      | 68.50                                      | 75.05                                  | 0.12  |
| Mean                   | 51.88  | 56.79                                      | 53.08                                      | 48.60                                      | –                                      | –     |
| CD (0.05)              |  |  |  |  |  |       |
| Month=2.34             |  |  |  |  |  |       |
| Treatment=1.65         |  |  |  |  |  |       |
| Month × treatment=4.68 |  |  |  |  |  |       |

**Table 7.** Average green and dry weight of shoots (t/ha) and moisture percentage of wicker willow (*Salix triandra* L.) before leaf fall on 15 September.

| Treatments                        | Green weight<br>(t/ha) | Dry weight<br>(t/ha) |
|-----------------------------------|------------------------|----------------------|
| T <sub>1</sub> (fresh plantation) | 16.7                   | 6.78                 |
| T <sub>2</sub> (2 yr old coppice) | 34.6                   | 14.20                |
| T <sub>3</sub> (3 yr old coppice) | 42.4                   | 18.24                |
| T <sub>4</sub> (4 yr old coppice) | 54.3                   | 23.21                |
| CD (0.05)                         | 0.73                   | 0.17                 |

The available nitrogen, phosphorus and potassium were found in the range of low, high and medium (Table 1). The concentration of available nitrogen was low due to the reasons that the short rotation forest species grow rapidly and bind great amount of nutrients in their biomass and also losses due to leaching, denitrification and ammonification of nitrogen is high. Phosphorus concentration increases due to the addition of litter fall which results into formation of polyhumic substances and coating of aluminium and iron ions thus the higher content of phosphorus in soil may be attributed to the higher content of organic matter in the soil which prevents the fixation of phosphorus ions. Potassium falls in the medium range due to the reason that potassium is always in the exchangeable form in the soil. So it indicates that nutrient status of the soil is important in short rotation forestry which is in confirmation with earlier reports (14).

The studies conducted on the growth of *Salix triandra* L. on various parameters revealed that sprouting vigor in T<sub>1</sub> (fresh plantation) was slow than the other treatments (Table 2). On 30 March, 100% sprouting was recorded in T<sub>3</sub> (three year old coppice) and T<sub>4</sub> (four year old coppice) stand. Whereas, in fresh and one year old coppice it was observed upto

**Table 8.** Average green/dry yield in (t/ha) of wicker willow (*Salix triandra* L.) after leaf fall on 15 November.

| Treatments                        | Green yield<br>(t/ha) | Dry yield<br>(t/ha) |
|-----------------------------------|-----------------------|---------------------|
| T <sub>1</sub> (fresh plantation) | 17.5                  | 7.85                |
| T <sub>2</sub> (2 yr old coppice) | 35.2                  | 15.76               |
| T <sub>3</sub> (3 yr old coppice) | 43.5                  | 19.00               |
| T <sub>4</sub> (4 yr old coppice) | 56.3                  | 25.10               |
| CD (0.05)                         | 1.41                  | 1.62                |



**Figure 1.** The rods recovered from wicker willow (*Salix triandra* L.) used as raw materials for manufacturing a variety of fancy items.

4 April. Statistically significant differences were found in  $T_1$  (fresh plantation) and  $T_2$  (two year old coppice) on 15, 20 and 25 March while  $T_3$  (three year old coppice) and  $T_4$  (four year old coppice) differed significantly on 15 March. The results indicate that more sprouting vigor in  $T_2$  (two year old coppice),  $T_3$  (three year old coppice) and  $T_4$  (four year old coppice) than  $T_1$  (fresh plantation) was due to the higher carbohydrate levels and well established root system in these coppice stands than the fresh cuttings  $T_1$  (fresh plantation), which diverted its reserve in root establishment. Also coppicing effect increases the hormonal changes in the remaining stumps, as willow sprouts from the auxillary buds of recent sprouts and enhances the sprouting due to the hormonal changes. These results are in agreement with earlier reports (15—17).

The number of shoots per coppice is presented in Table 3 which revealed that maximum shoots per coppice were recorded in  $T_4$  (7.66) on 15 April and minimum on 15 November (6.97) followed by  $T_3$ ,  $T_2$  and  $T_1$ . The mortality percentage was highest in  $T_4$

(9.00%) followed by  $T_3$  (5.68%),  $T_2$  (4.66%) and lowest in  $T_1$  (4.41%), in one growing season. So it can be concluded that both number of shoots and mortality percentage increased with increase in age of the coppice. Obviously due to the reason that the fast growing species needs complete over head light so with the increase in shoots the maximum shoots does not capture complete light, thus resulting lower photosynthetic efficiency which in turn accounted for reduction in net assimilation rate, so mortality percentage was increased. These results are in conformity with the earlier reports (2, 18—20).

*Salix triandra* L. showed three main periods of growth, namely 15 April to 15 June, 15 July to 15 August and 15 August to 15 November (Tables 4—6). The average shoot height average collar diameter and number of leaves increased significantly rapidly during the first two stages and then in third stage it increased with decreasing rate and finally becomes steep. The highest average mean increment in shoot height (55.7 cm), collar diameter (0.61 cm) and number

**Table 9.** Dimensions of basket willow of *Salix triandra* L. (wicker willow).

| Grade  | Required length (cm) | Observed length (cm) | Required diameter (cm) | Observed diameter (cm) |
|--------|----------------------|----------------------|------------------------|------------------------|
| Thin   | 80—160               | —                    | 0.2—0.4                | —                      |
| Medium | 180—280              | 162.90—187.30        | 0.5—1.1                | —                      |
| Thick  | 300—400              | —                    | 1.2—1.5                | 1.26—1.54              |

of leaves (22.45) was recorded from 15 June to 15 July. The average maximum shoot height (142 cm), collar diameter (1.12 cm) and number of leaves (6.79) was recorded in T<sub>2</sub> (two year old coppice). Statistically, the increase in average height, collar diameter, number of leaves differed significantly in T<sub>1</sub> (fresh plantation), T<sub>3</sub> (three year old coppice) and T<sub>4</sub> (four year old coppice) upto August while in T<sub>1</sub> (fresh plantation) it increased significantly upto 15 September. Maximum mean average height (187.64 cm), collar diameter (1.51 cm) and number of leaves (75.05) were recorded on 15 November and minimum value of average height, collar diameter and number of leaves on 15 April (20.15 cm), (0.10 cm) and (8.07), respectively. These values were attained in a period of one growing season which indicated the fast growth of this species. The shoot height, collar diameter and number of leaves were found to be minimum in T<sub>3</sub> (three year old coppice) and T<sub>4</sub> (four year old coppice) due to the reasons that the number of shoots were higher than that of other T<sub>1</sub> (fresh plantation) and T<sub>2</sub> (two year old coppice) and the nutrients get distributed in the more number of shoots. In T<sub>1</sub> (fresh plantation) the root system was not fully developed so maximum nutrients were used by the cutting for the development of root system. These results are in conformity with earlier findings (20—22).

The results indicated that maximum growth occurred during the warmer seasons i.e. June and July with rich precipitation (28.3 mm rainfall), and the species was found to invest more photosynthetic products to above ground parts during the early growth which ultimately facilitated plants to capitalize more light and allow fast above ground growth (12, 23).

Harvesting index is reported to be one of the important factors for the management of coppice stand. The harvesting was observed best when the color of the lower leaves started to change from green to brown and finally turned yellowish. While harvest-

ing time started when complete or greater than 75% of leaf fall occur/shedding took place i.e. during dormant period. It is obviously due to the reason that when days become shorter and nights longer it indicate the harvesting stage. So with the decrease in photoperiod, temperature and number of leaves, photosynthesis decreased and content of abscisic acid increases, which indicated dormant period was best time of harvesting of *Salix triandra* L. These results are in agreement with those recorded earlier (2, 24, 25). The relevant data on green and dry weight of shoots before and after leaf fall are presented in Tables 7 and 8 which reveal that with the increase in age of the coppice the green and dry weight increased significantly. The maximum green weight of shoots was recorded in T<sub>4</sub> (four year old coppice) (54.3 t/ha) and lowest in T<sub>1</sub> (fresh plantation) (16.7 t/ha), while dry weight values recorded after leaf fall for the treatments were subsequently maximum in T<sub>4</sub> (four year old coppice) (23.21 t/ha) and lowest in T<sub>1</sub> (fresh plantation) (6.78 t/ha). The maximum shoot weight in T<sub>4</sub> (four year old coppice) was obviously due to the maximum number of shoots than other treatments. The variation in yield before leaf fall and after leaf fall was due to the reason of continued photosynthesis. The green and dry yield reached maximum when complete leaf fall took place i.e. when photosynthesis ceases. The moisture per cent calculated did not differ significantly among the treatments because in all the treatments growth was of one year and moisture per cent ranged between 56.32 to 55.14. Due to the obvious reasons that one year shoots grows fast to capture light and LAI reaches maximum which increases the rate of photosynthesis; these results were in agreement with the earlier findings (22, 26).

Processing is important for storage of raw material. After boiling for 10—12 hours with hies or rods debarked by fastening the bark in the two wooden sticks embedded one-third in the soil. Then dried for

**Table 10.** Average rod (withies) diameter and length after processing of wicker willow (*Salix triandra* L.) (cm). \*Diameter recorded from thick end.

| Treatments                        | *Diameter after debarking (cm) | Length after debarking (cm) |
|-----------------------------------|--------------------------------|-----------------------------|
| T <sub>1</sub> (fresh plantation) | 1.52                           | 186.5                       |
| T <sub>2</sub> (2 yr old coppice) | 1.54                           | 187.3                       |
| T <sub>3</sub> (3 yr old coppice) | 1.34                           | 175.74                      |
| T <sub>4</sub> (4 yr old coppice) | 1.26                           | 162.90                      |
| CD (0.05)                         | NS                             | 6.49                        |

a week's time under sun light. Boiling softens the bark and bark tannins was absorbed into stem wood which gives buff tint to rod and make it water proof and increases its durability, and prevents the raw material from deterioration, increases strength and quality. The sun drying is found to be most cost effective and the rods dried 10 times slower with bark than debarked because maximum moisture per cent is present in bark (27).

Raw materials after processing were graded on the basis of rod length and diameter. The average rod length varied from 162.9 to 187.3 cm and average diameter 1.26—1.54 cm (Table 9). It indicates that average length falls in the medium and diameter in the thick grade—these dimensions are in agreement with the dimensions of standards (5).

Average rod length and diameter after processing were measured which revealed that the maximum length and diameter were found in T<sub>2</sub> (two year old coppice) (187.30 and 1.54 cm) followed by T<sub>1</sub> (fresh plantation) (186.50 and 1.52 cm), T<sub>3</sub> (three year old coppice) (175.74 and 1.34 cm) and T<sub>4</sub> (four year old coppice) (162.90 and 1.26 cm) (Table 10). Statistical analysis revealed that length of the rods differed significantly among the treatments, while collar diameter was not found to be significant among the treatments. The rod length and collar diameter were found to be less than that recorded before processing due to the reasons that during processing the bark and succulent tips were removed. The bark and tips removed contributed nearly 25% of its total weight. These results were in confirmation with the earlier findings (28).

### Conclusion

Results from the study showed that the soil pH

was neutral to acidic, EC medium, organic carbon high with available NPK, low, high and medium respectively. Sprouting vigor was maximum in T<sub>3</sub> (three year old coppice) and T<sub>4</sub> (four year old coppice) which commenced and completed earlier than T<sub>2</sub> (two year old coppice) and T<sub>1</sub> (fresh plantation). The number of shoots per coppice was the deciding factor in contributing yield. However, the bark contributed nearly 25% of the total weight. Dimension of processed rods falls in medium category in length and thick category in diameter.

### References

1. Alvium R., E. W. Hewett and P. F. Savnders. 1976. Seasonal variation in the hormone content of willow, change in abscisic acid content and cytokinin activity in the xylem sap. *Pl. Physiol.* 57 : 474—476.
2. ASAE (American Society of Agricultural Engineers). 1994. Moisture measurements—forages, ASAE standards S 358.2 Dec 93.
3. Blake T. J. 1983. Coppice systems for short-rotation intensive forestry. The influence of cultural, seasonal and plant factors. *Australian For. Res.* 13 : 279—291.
4. Bullard M. J., S. J. Mustil, S. D. McMelian, B. M. Nixon and C. P. Brilit. 2002. Yield improvements through modification of planting density and harvest frequency in short rotation coppice *Salix* spp. Yield response in two morphologically diverse varieties. *Biom. and Bio-En.* 22 : 15—25.
5. Cannell M. G. R. 2004. Short rotation forestry for biomass production. Pages 874—875 in B. Jaffery, E. Julain, Y. A. Jhon, editors. *Encyclopedia of forest-Sciences*, volume 2.
6. Cannell M. G. R., L. J. Sheppard and R. Milne. 1988. Light use efficacy and woody biomass production of poplar and willow. *Forestry* 61 : 125—136.
7. Christersson L. 2005. Plant physiological aspects of woody biomass production for energy purposes. Short rotation forestry for industrial and rural development. Pages 1—7 in K. S. Verma, D. K. Khurana and L. Christersson, editors. *ISTS*, Nauni, Solan, India.
8. Crouch R. J. and M. N. Honeyman. 1986. The relative salt tolerance of willow cuttings. *J. Soil Conserv., New-South-Wales* 42 : 103—104.
9. Decei I. 1975. Amount and rate of weight loss in willow osiers, *Studii-Si-Cercetari*, Institutu-de-Cercetari-Si- Amenajari-Silvice, I-Silvicultura 33 : 111—118.
10. Ericsson T. 1981. Growth and nutrition of three *Salix* clones in low conductivity solutions. *Physiologia-Plantarum* 52 : 239—244.
11. Gigler J. K., W. K. P. Loon Van, J. V. Berg Vanden, C. Sonneveld and G. Meerdink. 2000. Natural wind drying of willow stems. *Biom. and Bioen.* 19 : 153—163.
12. Hytonen J. 1996. Biomass production and nutrition of short rotation plantations. *The Finnish For. Res. Inst.*

- Res. Paper, pp. 556—561.
13. Jackson M. L. 1973. Soil chemical analysis, 2nd edition. Prentice Hall, New Delhi, India.
  14. Kauppi A., M. Kiviniitty, C. Tigerstedt and A. N. Ferm. 1987. Number and location of basal bunds in *Alnus incana* and *Salix Aquatica*. Poster-preparation in IEA/bioenergy meeting in Qulu. Finland, pp. 24—29.
  15. Krussman G. 1978. Cultivated broad leaved trees and shrubs, volume 3. Timber Press, Portland, Oregon.
  16. Nordberg S. and K. Vertaileva. Pajin viijelykseen Jasen edellytyksiin ulkomailla ja suomessa. Referat 1928. Die Weidenkultur und-ihre vovausset-zungen in Ausland und Suomi. *Silva Fennica* 9 : 1—63.
  17. Olsen S. R., W. Cole, F. S. Watanable and L. A. Dean. 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate Pages 1044—1046 in C. A. Black, editor. *Methods of soil analysis*. American Soc. Agron. Madison, USA.
  18. Perttu K., H. Eckersten, P. J. Kowalik and L. O. Nilsson. 1984. Modelling potential of energy forest production. Pages 467—480 in K. Perttu, editor. *Ecology and management of forest biomass production system*, Report 15. Swedish Univ. Agric. Sci. Dep. Ecol. and Environm. Res.
  19. Romero-Ablos M. I. 1988. Towards development of the Chilean basket willow sector <http://www.Fao.org/docresp/008/a0026e/a0026e10.htm>. papers and willows.
  20. Savill P. S. 2004. Temperate broad leaved deciduous forest. Pages 1398—1403 in B. Jaffery, E. Julain, Y. A. Jhon, editors. *Encyclopedia of forest sciences*, volume 3.
  21. Sennerby-Forsse L., H. Johansson. 1990. *Energiskog Handboki Praktisk Odling*. Uppsala Solar.
  22. Smallukas D. and R. Noreika. 2005. Morphological characteristics and evaluation of productivity of *Salix triandra* L. clones selected for short-rotation coppice. *Biologija* 3 : 36—40.
  23. Subbiah B. V. and G. L. Asija. 1956. A rapid procedure for estimation of available nitrogen in soils. *Curr. Sci.* 25 : 259—260.
  24. Szczukawski S., M. Stolorski, J. Twarkowski, J. Przyborowski and A. Klasa. 2005. Productivity of willow coppice plants grown in short rotations. *Pl. Soil Environ.* 51 : 423—430.
  25. Taylor J. S., T. J. Blake and R. P. Pharis. 1982. The role of plant hormones and carbohydrates in the growth and survival of coppiced, *Eucalyptus* seedlings. *Physiol. Pl.* 44 : 421—430.
  26. Tolonen R. 1988. Early dynamics and height growth modelling of coppiced *Betula pubescens* stands (in Finnish). Master thesis, Dep. of Biophys., Univ. Qulu.
  27. Wadoo M. S. 2005. Veer—the multipurpose tree. Kashmir Observer [http://documents%20settings/administrator/desktop/Kashmir % 20 observation 1/28 2005](http://documents%20settings/administrator/desktop/Kashmir%20observation%201/282005).
  28. Walkley A. and C. A. Black. 1934. An examination of the Degtjareff method for determination of soil organic mater and proposed modification of the chromic acid titration method. *Soil Sci.* 37 : 29—38.